

B 146191


JATE
EGYETEMI GYŰJTEMÉNY
FELYBEN
OLVASHATÓ

PLANT CELL BIOLOGY AND
DEVELOPMENT

EDITED BY

M. KEDVES


B




Wohles

PROPERTY OF
DEPARTMENT OF
COMMERCE

U.S. DEPARTMENT OF
COMMERCE
BUREAU OF
CENSUS



B 146191



**HELYBEN
OLVASHATÓ**

Plant Cell Biology and Development

5

Editor: M. Kedves

Financial support

by the Foundation for the Science of the South Hungarian Plain,

by the Scientific and High-educational Committee of the Local Government of

Szeged,

by the J. A. University and the Faculty of Sciences,

by the Grant OTKA 1/3, 104

JATE Egyetemi Könyvtár



J000092248



B. 146191

This volume is scientific exchange matter and no commercial

HU ISSN 0866-5443

1994
Szeged

Contents

Preface	7
To the Tenth Anniversary of the Discovery of Quasicrystals KEDVES, M.	9
1. LM Investigations of different stained fossil <i>Botryococcus</i> colonies VÉR, A.	11
2. Dorog type (Middle Eocene) spore-pollen assemblage in the Socka Beds of Slovenia I. JELEN, B., KEDVES, M., SKABERNE, D., BREZIGAR, A., BUSER, S., CIMERMAN, F., DROBNE, K., MONOSTORI, M., PAVLOVEC, R., and PAVŠIČ, J.	20
3. Plant microfossils from the Upper Cretaceous and Lower Tertiary layers of Northern Spain I. KEDVES, M.	29
4. Consideraciones en torno a las asociaciones esporopólicas del Cretácico Superior del Cerro de la Mesa (Guadalix de la Sierra, Madrid) ALVAREZ RAMIS, C., KEDVES, M. y FERNÁNDEZ MARRÓN, T.	42
5. TEM study of the "infected" pollen grains of <i>Thalictrum flavum</i> L. with <i>Gliocladium roseum</i> (LINK) BAINIER KEDVES M. and ROJIK, I.	52
6. Buckminsterfullerene-like biopolymer units from the exine of <i>Thalictrum flavum</i> L. KEDVES, M. and ROJIK, I.	58
7. Incomplete and non-fivefold rotation of the basic biopolymer unit of the exine of <i>Pinus griffithii</i> McCLELL KEDVES, M., TÓTH, A. and GOTTL, E.	67
8. Premiers résultats du système de biopolymère stabilisateur du squelette quasi-cristalloïde de l'exine KEDVES, M. et TÓTH, A.	79
9. List of publication of M. KEDVES II. GOTTL, E.	87
Chronicle KEDVES, M. and GÁSPÁR, I.	101

Contributors

Concepcion ALVAREZ RAMIS

Department of Paleontology. Faculty of Geology. U. C. M. Madrid, Spain,
University Professor.

Aleksander BREZIGAR

Geological Survey of Ljubljana. Institute for Geology, Geotechnics and
Geophysics, Ljubljana, Slovenia. Head of Department of Geophysics.

Stanko BUSER

Department of Geology and Paleontology. University of Ljubljana. Ljubljana,
Slovenia. University Professor.

Franci CIMERMAN

Paleontological Institute Ivan Rakovec, ZRC SAZU. Ljubljana, Slovenia,
Research Adviser.

Katica DROBNE

Paleontological Institute Ivan Rakovec, ZRC SAZU. Ljubljana, Slovenia. Head
of Paleontological Institute.

Maria Teresa FERNÁNDEZ MARRÓN

U. E. I. of Paleontology. Institute of Economic Geology (C. S. I. C. – U. C. M.)
Madrid, Spain, Researcher.

István GÁSPÁR

Cell Biological and Evolutionary Micropaleontological Laboratory, University
Student.

Elza GOTTL

Cell Biological and Evolutionary Micropaleontological Laboratory.
Grammar-school Teacher.

Bogomir JELEN

Geological Survey of Ljubljana. Institute for Geology, Geotechnics and
Geophysics, Ljubljana, Slovenia. Senior Research Associate.

Miklós KEDVES

Cell Biological and Evolutionary Micropaleontological Laboratory. Research
Councilor. Honorary Professor.

Miklós MONOSTORI

Department of Paleontology, E. L. University. Budapest. Associate Professor.

Rajko PAVLOVEC

Department of Geology and Paleontology. University of Ljubljana. Ljubljana,
Slovenia. University Professor.

Jernej PAVŠIČ

Department of Geology and Paleontology. University of Ljubljana. Ljubljana,
Slovenia. University Professor.

Imre ROJIK

Faculty of Sciences of the J. A. University. Laboratory of Electron-microscopy.
Research Ass. Professor.

Drago SKABERNE

Geological Survey of Ljubljana. Institute for Geology, Geotechnics and Geophysics, Ljubljana, Slovenia. Senior Research Associate.

Anita TÓTH

Cell Biological and Evolutionary Micropaleontological Laboratory. University Student.

Annamária VÉR

Cell Biological and Evolutionary Micropaleontological Laboratory. University Student.

Preface

This number of the laboratory publications is dedicated to the Tenth Anniversary of the Discovery of the Quasicrystals. I hope, that the presented papers herein represents well the aim of our effort, namely the following:

- i. The inter- and/or multidisciplinary researches in our fields.
- ii. The adaptation of the methods of other fields of researches.
- iii. In the same time to continue so-called classical researches with classical methods, but with the complementation of the newest molecular or biophysical concepts.

The discovery of the quasicrystals and the fullerenes are very important to the research programs of our laboratory. The last years, and it seems in the future we will carry out several experimental studies with the concept of the quasi-crystalloid skeleton of the plant cell wall and/or of the cytoskeleton too.


This number was published by the financial support of several institutions. I would like to express my sincere thanks to the following persons and institutions:

to Prof. Dr. GY. TELEGDY member of the Hungarian Academy of Sciences,
to the Foundation for the Science of the South Hungarian Plain,
to the Scientifical and High-educational Committee of the Local Government of Szeged,
to Prof. Dr. I. DÉKÁNY scientifical vice-rector of the J. A. University, and Prof. Dr. R. MÉSZÁROS, Dean of the Faculty of Science,
to the Foundation of the Grant OTKA 1/3–104.

Szeged, 25 February 1994.

M. KEDVES
head of the laboratory

THE TENTH ANNIVERSARY OF THE DISCOVERY OF QUASICRYSTALS



Group of scientists with three of the discoverers of quasicrystals, Dan SHECHTMAN (1), Denis GRATIAS (2) and John W. CAHN (3). The picture was taken from a NIST publication (1988), and it does not include Professor Ilan BLECH. The picture was made available for us by courtesy of Dan SHECHTMAN, the reproduction was made by Dr. I. BAGI.

TO THE TENTH ANNIVERSARY OF THE DISCOVERY OF QUASICRYSTALS

M. KEDVES

Cell Biological and Evolutionary Micropaleontological Laboratory of the Department of Botany of the J. A. University, H-6701, P. O. Box 657, Szeged, Hungary

It was ten years ago that SHECHTMAN, BLECH, GRATIAS and CAHN published rapidly cooled AlMn alloy that gave diffraction patterns with full icosahedral symmetry. This discovery indicated an extremely intensive research in several fields of Crystallography, Physics, Mathematics, etc.

Several handbooks have been published and synopsis have been organized on this problem and with its inter- or better say multidisciplinary character and connections. The number of the papers published on the new results is enormous, it is extremely difficult to enumerate only the most important steps in this field of research during the last few years. Some selected examples may be mentioned as follows.

1989 – Symmetry of structure. An interdisciplinary Symposium (Abstracts in two volumes, edited by GY. DARVAS and D. NAGY).

1990 – Quasicrystals, Network, and Molecules of Fivefold Symmetry. VCH Publishers, Inc. New York, I. HARGITAI editor; contributors: A. L. MACKAY, M. SENECHAL, R. Z. SAGDEEV and G. M. ZASLAVSKY, PENG JU LIN and L. A. BURSILL, F. DÉNOYER, G. HEGER and M. LAMBERT, R. A. DUNLAP, D. W. LAWTHORP and V. SRINIVAS, J.-P. ALLOUCHE and O. SALON, E. J. W. WHITTAKER and R. M. WHITTAKER, M. BAAKE, P. KRAMER, M. SCHLOTTMANN and D. ZEIDLER, P. McMULLEN, H.-U. NISSEN, P. STAMPFLI, P. G. MEZEY, D., J. KLEIN and T. G. SCMALZ, A. LÉGER, L. D'HENDECOURT, L. VERSTRAETE and W. SCHMIDT, E. BRENDSDAL, S. J. CYVIN, B. N. CYVIN, J. BRUNVOLL, D. J. KLEIN and W. A. SEITZ, V. ELSE, E. BRENDSDAL, S. J. CYVIN, J. BRUNVOLL, B. N. CYVIN and D. J. KLEIN, E. BRENDSDAL, J. BRUNVOLL, B. N. CYVIN and S. J. CYVIN, D. KUCK.

1991 – Topics on contemporary Crystallography and Quasicrystals, Special Issue, Periodico di Mineralogia. Edited by L. LORETO and M. RONCHETTI; contributors: M. V. JARIC, H.-U. NISSEN and C. BEELI, S. MERLINO, M. EMMER, R. PENROSE, L. H. TANG, M. RONCHETTI, C. ANTONIONE, L. BATTAZZATI, M. CALLERI and F. MARINO, A. L. MACKAY, L. LORETO, R. FARINATO and F. PAPPALARDO, A. L. LOEB.

M. RONCHETTI compiled the bibliography of Quasicrystals.

1991 – Symmetry and Topology in Evolution, edited by B. LUKÁCS, SZ. BÉRCZI, I. MOLNÁR and G. PAÁL.

1992 – Szimmetria – Aszimmetria, edited by T. BALOGH, MTA SZAB.



After these examples, let me make a short list of the most important results of our laboratory concerning quasicrystals in the field of Cell Biology.

- 1988 – Discovery of the quasicrystalloide biopolymer structure in living system on partially degraded exine of *Pinus griffithii* McCLELL.
- 1989 – A modified MARKHAM rotation method was elaborated to investigate the symmetries of the basic quasi-crystalloid skeleton of the plant cell wall. Highly organized biopolymer structures were described with quasi-crystalloid basic units. A comprehensive model of the biopolymer organization of the sporoderm was established. Three major degrees were distinguished at the highly organized biopolymer system of the sporoderm.
- 1990 – The first TEM picture was published about the PENROSE unit-like biopolymer unit from the partially degraded exine of *Pinus griffithii* McCLELL obtained with the secondary rotation method. The first data about the biopolymer structure of partially degraded wall of the spores of the genus *Selaginella*.
- 1991 – The first observations of the biopolymer structure of the intine. Biopolymer units of hexagonal symmetry were described. Data about the superficial molecular system of the pollen wall. Highly organized globular biopolymer units from the partially degraded and fragmented wall of *Botryococcus braunii* KÜTZ. extracted from the Hungarian oil shale were published. These units can be modelled with fullerenes, which are quasi-equivalent biopolymer structures. Quasi-crystalloid biopolymer structures were described from explosive dangerous Jurassic coal pulver. Quasi-crystalloid biopolymer structures were published from partially degraded and fragmented sclereids of *Armeniaca vulgaris* LAM. Three-dimensional modelling of the biopolymer skeleton of the plant cell wall.
- 1992 – Secondary alterations in the biopolymer structure of the exine as a consequence of the X-ray effect. The first data about the stabilizing system of the quasi-crystalloid biopolymer skeleton of the sporoderm.
- 1993 – “Negative” quasi-crystalloid biopolymer network was described from the partially degraded exospore of *Equisetum arvense* L. The radial rotation method was introduced with the alterations of the outer and inner rotation areas as a new two-dimensional method.

In resumé, the inter- or multidisciplinary principles can result several new aspects in the so-called “classic” scientific fields. The new research fields of Crystallography have opened new aspects and prospects in Cytology, and in the investigations of fossil organic material. At the recent investigation material, not only the biopolymers of the plant cell wall can be investigated with the new research concepts and methods, but also the cytoskeleton and all kinds of organelles of living systems. The discovery of quasicrystals was an extremely important contribution to this new research on recent and fossil plant organelles.

1. LM INVESTIGATIONS OF DIFFERENT STAINED FOSSIL BOTRYOCOCCUS COLONIES

A. VÉR

Cell Biological and Evolutionary Micropaleontological Laboratory of the Department of Botany of the J. A. University, H-6701, P. O. Box 657, Szeged, Hungary

Abstract

Colonies of the *Botryococcus braunii* KÜTZ. algae extracted from the Upper Tertiary oil shale of Pula (Hungary) were investigated with the light-microscope. Stained, non-stained and solvated colonies were the subjects of these researches. The stain acceptance is suitable for researches of this kind, too, to establish the degrees of maturity of the organic components of the colonies.

Key words: *Botryococcus braunii*, oil shale, Neogene, stain acceptance, Pula, Hungary.

Introduction

The basic morphology of the genus of the *Botryococcus* KÜTZ. is extremely peculiar and isolated in the Plant Kingdom. This is the reason why the taxonomic position has been widely discussed and changed. This genus was ranged with the *Heterocontae* (Kiss, 1939) respectively within the *Chlorophyceae* WILLE (in ENGLER and PRANTL, 1910). Morphological statements by FREMY and DANGEARD (1938) and CHADEF AUD and EMBERGER (1960), COMBAZ (in DURAND, 1980) and GLIKSON, LINDSAY and SAXBY (1989). Resuming the basic morphology is as follows (Plate 1.1.).

1. The colonies are enveloped in mucilage (= Schleimmasse = mucilagineuse = une enveloppe gelatineuse hyaline = hydrocarbon matrix).
2. The differentiations of the cellular components of the colonies are: pedunculus, cupula. The individuum are in pair in the cupula and are forming one "polipier" sensu CHADEF AUD and EMBERGER (1960).
3. COMBAZ (1980) (Cf. CHADEF AUD and EMBERGER, 1960), published a scheme about the organelles of the protoplasm (nucleus, amidon, oil drops, plastis). An important component is the cuticle cellular. The cuticle of the mother cell is a part of the basal region of the "polipier" (Plate 1.1.). The morphology of the pedunculus is very characteristic. (Plate 1.1., 1.2.)

TEM data about fossil *Botryococcus* colonies from the oil shale were published by KEDVES (1983) and GLIKSON, LINDSAY and SAXBY (1989). The oil reservoir function of the holes in mucilage and in the more or less compact wall substance of the fossil colonies was pointed out (KEDVES, 1983). Biopolymer organization (KEDVES, 1988)

and thin layer chromatography was first described in two papers from the locality of Pula (KEDVES, 1986a, b).

Concerning the ecology of the *Botryococcus braunii* KÜTZ. algae the following can be pointed out:

- i. Occurrence in eutrophic fresh water or in humid soil (WILLE, in ENGLER and PRANTL, 1910) in Europe, North America and Africa. These colonies can also be present in salt lakes or in the water of marine lagoons.
- ii. Stenotherm species after the paper of KISS (1939). The geological distribution of this kind of algae is extremely large. JARZEN (1978) wrote the following, p. 32:
"Botryococcus KÜTZING (Pl. 1, fig. 3) is a colonial green algae, whose colonies form irregular globose masses encasted in a heavy, often dark, cohered mucilage. TRAVERSE (1955) have reviewed the fossil occurrences of the genus and notes that the fossil record probably extends back at least to the Ordovician." (Cf. NARAYANA RAO and MISRA, 1949).
"The hydrocarbon secreting alga *Botryococcus* has been identified in organic remains of sediments ranging from Precambrian to Recent," (GLIKSON et al., 1989, p. 595).

Taking into consideration several results the chemical compositions of the fossil forms is determined by the following factors:

1. The basic composition of the different cellular elements of the colonies.
2. The ontogenetical state of the algae.
3. The molecular alterations in the wall also during the life.
4. The fossilization of the algae.
5. Taphonomical processes.
6. The preservation in the sediments.

The aim of this research is complex. By the LM method and the stain acceptance to get information, about the maturity of the different part of the colonies (Cf. POTONIE and REHNELT, 1971). The advantage of the light-microscopy combined with cytochemical methods is that a mass of data can be obtained. The statistical evaluation of the great number of information is important from the point of view of the Ontogenesis, Ecology, and Taphonomy.

Materials and Methods

The material of investigation is an average sample from the oil shale deposits of Pula (Hungary, Transdanubia). The oil shale of Transdanubia (Hungary) was first discussed by JÁMBOR and SOLTÍ (1975). Following JÁMBOR (1980) there are peculiar geological and paleoenvironmental conditions during the formation of these volcanic lakes. The water of the lakes is oligohaline water. The Geology (Cf. JÁMBOR, 1980, HETÉNYI, 1985, JÁMBOR and SOLTÍ, 1975, SOLTÍ, 1981), Ecology (Cf. NAGY, 1976, HAJÓS, 1976) and Geochemistry (HETÉNYI, 1985, 1987–1988, HETÉNYI and PÁPAY, 1986, ARATÓ and BELLA, 1976) of the oil shale was the subject of several investigations. The most important statements are as follows:

1. The investigated oil shale is a volcanic lake type. (JÁMBOR and SOLTÍ, 1975, JÁMBOR, 1980).
2. There was a geysirite activity during the sedimentation of the

alginite (SOLTI, 1981). 3. The water was oligohaline 0.3‰, indicated by the diatoms published by HAJÓS (1976). 4. The clima was warm-temperate (annual average temperature: 10–12 °C), following HAJÓS (1976). E. NAGY (1976) described the pollen grains of Mediterranean conifers. 5. The water depth was some meters only based on the benthonical algae (Cf. HAJÓS, 1976). 6. The riparian Woodland resulted a peculiar local clima (E. NAGY, 1976). 7. The above mentioned conditions resulted the water flowering with the *Botryococcus braunii* forming the alginite.


The pulverized material was mixed with water. After stirring the inorganic components settled down and the organic fraction was separated, and newly dried. The investigation was prepared as follows. 1. Intact colonies (after separations mounted in glycerin-jelly hydrated at 39.6%). 2. Colonies partially dissolved with methanol. 3. To get information about the alterations or maturity of the colonies the following stains were used: 3. 1. BISMARCK Brown (vesuvin) 1% dissolved in ethanol 30%. 3. 2. BISMARCK Brown (vesuvin) 1% dissolved in ethanol 70%. 3. 3. Chrysoidin 2% dissolved in ethanol 40%. 3. 4. Safranin T 2% dissolved in ethanol 50%. 3. 5. Eosin 1% dissolved in distilled water. 3. 6. Fe-Haematoxylin (after EHRLICH). 3. 7. Malachite Green 1% dissolved in distilled water. 3. 8. Methylene Blue 1% dissolved in distilled water. 3. 9. Toluidine Blue 0.2% dissolved in distilled water.

The pictures were taken with an objective of oil immersion; Carl Zeiss Jena, GF Planachromat HI 100x/1.25∞/0.17–A, except, Plate 1.1.

Results

The characteristic morphology with the different cellular organelles is illustrated in Plate 1.1. The pictures were taken from colonies stained with Toluidine Blue, with an objective Carl Zeiss Jena, GF Planachromat 40x/0.65∞/0.17–A. Plate 1.2. illustrates examples about the degradation and the kerogen accumulation. The description of the detailed results is given in the order under the title Materials and Methods.


1. Intact colonies (Plate 1.2., figs. 3,4, and 13,14). It is a relationship between the preservation stage and geochemical maturity of the colonies. The accumulation of kerogen may be indicator of the “maturity level.” In the case of kerogen accumulation, the colonies are at least partially degraded (Plate 1.2., figs. 13,14). The ontogenetically primitive colonies are less degraded in contrast to the completely developed ones. The presence of the green granules is also an indicator of the maturity level. On the basis of the data of the material of Pula the following can be distinguished. 1. 1. Clear greenish-yellow, semi-transparent colonies without kerogen accumulation and/or green granules. 1. 2. Clear greenish-yellow to brownish-yellow colonies with green granules. Colonies of this kind may be degraded to some level. 1. 3. Partially degraded colonies containing kerogens (brown particles), and green granules. 1. 4. The advanced degradation process is well shown with relatively high quantity of kerogens and green granules. 2. Colonies partially dissolved with methanol (Plate 1.2., figs. 7,8, and 9,10). The preliminary investigations did not result in characteristic alterations. The colonies are more transparent in contrast to the intact ones only. Further experiments are necessary with this solvent. 3. Stained colonies (Plate 1.1., 1.2., figs. 1,2,5,6, 11,12, 15–21). In comparison to the intact colonies the following characteristic



features were investigated by the stained material: the green granules; the brown particles of the kerogen; the stain acceptance of the mucilage; the different particles of the cupula and the pedunculus. Particular attention was paid to the degradation surfaces. 3.1. and 3.2. BISMARCK Brown (vesuvin) 1% dissolved in ethanol 30% and 70% (Plate 1.2., figs. 1,2,11,12). No significant differences were found between the two kinds of dissolved stains. The colonies have not accepted this stain but the solvent effect of the ethanol was observed which is similar to those of the methanol (2.) 3.3. Chrysoidin 2% dissolved in ethanol 40% (Plate 1.2., fig. 20). The inner regions of the wall respectively the cupula and the pedunculus accepted this stain but only at the well preserved colonies. The colour is orange occasionally dark orange, or yellow. Worth of mentioning is the unstained part of the wall. The granules are stained at the well preserved colonies in a middle per cent. The granules of the degraded colonies have not accepted the stain or only to a very limited measure. The colour of the completely damaged colonies is yellow. No colour change was observed at the brown particles and the mucilage, in contrast to the non-stained colonies. 3.4. Safranin T 2% dissolved in ethanol 50% (not illustrated). The not degraded wall have accepted the stain only in a very poor measure. The degradations surfaces much better accepted the stain. The granules accepted occasionally the stain. The mucilage well accepted the stain in this way, its colour is red. The pedunculus is in general yellow, in this way this is the original colour. But the cupula have accepted partially the stain, the colour may be yellow or red. 3.5. Eosin 1% dissolved in distilled water (Plate 1.2., figs. 15,16). The wall accepted the stain, the colour is orange. The kerogen (the brown particles) is as at the non stained material. At some degraded colonies the colour of the cupula is orange. But in general it seems that the cupula and the pedunculus have not accepted the stain. The colour is as originally. The mucilage also accepted this stain. 3.6. Fe-Haematoxylin (after EHRLICH) (not illustrated). The slide colour changed; after a certain time to yellow. The colour of the wall is grey or greyish-blue. The cupula accepts the stain much more than the pedunculus. The colour of the pedunculus is originally yellow. The granules are green or occasionally bluish-violet. The colour of the kerogen has not changed; brown. 3.7. Malachite Green 1% dissolved in distilled water (not illustrated). The colour of the pedunculus at the well preserved colonies is yellow or transparent. At the partially degraded colonies the colour of the pedunculus is yellow. The pedunculus accepts the stain at the degraded specimens so the colour is green. The cupula generally accepts the stain. But exceptionally, when the pedunculus has not accepted the stain, the colour is yellowish-green. The mucilage accepts the stain very well, its colour is dark green. Kerogen accepts the stain. 3.8. Methylene Blue 1% dissolved in distilled water (Plate 1.2., figs. 17,18). The wall does not accept the stain in contrast to the granules. The outer part of the cupula is blue, the inner part is yellow. The pedunculus is yellow only rarely accepts the stain. The kerogen accepts the stain. 3.9. Toluidine Blue 0.2% dissolved in distilled water (Plate 1.1., 1.2., figs. 19,21).

◀ Plate 1.1

Botryococcus braunii KÜTZING from Alginite of Pula. General aspect from well preserved colonies from the Upper Neogene of Pula. Coloured with Toluidine Blue. The different cellular organelles are indicated. 1200 x.



The colonies accept the stain, the colour is green. This indicates the accumulation of the aromatic lignin derivates. The granules rarely accept the stain, at the well preserved colonies. At the degraded specimens the granules have been colored. The mucilage accepted the stain, colour: violet. The cupula is at the well preserved colonies green or greenish-yellow. The pedunculus is mostly violet. The colour of the degraded colonies and the kerogen is also violet.

Discussion and Conclusions

In consequence of the polycondensation and polymerisation during the diagenesis the kerogen is three dimensional network of organic molecules. The chemical composition has been investigated by several laboratories. From the very extensive data of a mass of publications the following citations will be pointed out.

DOUGLAS, EGLINGTON and MAXWELL (1969), p. 569: "The hydrocarbon content of coorongite, a Recent rubbery deposit derived from the alga *Botryococcus braunii*, has been investigated by infrared spectrometry, gas chromatography and combined gas chromatography-mass spectrometry."

ANDERS and ROBINSON (1971), p. 661: "Fifty-two cyclic alkanes, isolated from the bitumen of Green River shale, were analyzed by mass spectrometry."

The acceptance of the stains well indicates the heterogeneity of the colonies. Namely the colonies of different "maturity" and degradation level differentially accept the stains. The differences in the degree of the maturity well shown at the coloration with the Chrysoidin. The differences in the degradation level were demonstrated with the coloration of the Toluidine Blue, Safranin T and Malachite Green. Less expressed are the following stains: Chrysoidin and Eosin. Kerogen well accepted Safranin T, Malachite Green, Methylene Blue and Toluidine Blue. This fact indicates the presence of the aromatic derivates. The pedunculus accept less the stains in contrast

◀ Plate 1.2.

- 1-21. *Botryococcus braunii* KÜTZING from Alginite of Pula. 1000 x.
- 1,2. Well preserved colony coloured with Vesuvin.
- 3,4. Unstained slightly degraded colony with green granules.
- 5,6. Partially degraded colony stained with Eosin.
- 7,8. Completely degraded unstained colony, dissolved with methanol.
- 9,10. Kerogen accumulation at the basical part of the colony. Unstained material dissolved with methanol.
- 11,12. High measure of kerogen accumulation stained with Vesuvin (dissolved in ethanol 30%).
- 13,14. Unstained colony. The accumulation of the kerogen extends near to all parts of the colony. Unstained material.
- 15,16. The kerogen accumulation is at basis of the colonies. Stained with Eosin.
- 17,18. Well shown is the mucilage coloured with Methylene Blue.
19. The peculiar morphology of the pedunculus. The granules are coloured with Toluidine Blue.
20. Well preserved colony stained with Chrysoidin.
21. The granules coloured with Toluidine Blue.

C = cupula, M = mucilage, P = pedunculus.

to the cupula. Chrysoidin and Toluidine Blue coloured in an important measure the pedunculus. Malachite Green and Methylene Blue at the degraded colonies have coloured the pedunculus. It is necessary to point out that Toluidine Blue coloured the colonies in a different way. The colour of the cupula at well preserved colonies is green. The colour of the degraded cupules is violet. Methylene Blue coloured the outer part of the cupula. (Cf. CHADEFAUD and EMBERGER, 1960). The green granules have not totally accepted the stain. The stain acceptance was moderated at Chrysoidin, Toluidine Blue and Methylene Blue. In a very poor measure Safranin T, and Fe-Haematoxylin have coloured the green granules.

Resuming this was useful for several points of views. But it is necessary to develop this methodical concept, too. For this it is necessary to investigate the mechanism of the different coloration.

Acknowledgements

This work was supported by the grant OTKA 1/3, 104. The author is deeply indebted to Research Ass. Prof. Dr. M. HETÉNYI and to Prof. Dr. M. KEDVES for critically reading the manuscript.

References

- ARATÓ, Á. és BELLA, M. (1976): A pulai és gércei olajpala technológiai és kémiai vizsgálata. – MÁFI évi jel. az 1974. évről, 289–301.
- ANDERS, D. E. and ROBINSON, W. E. (1971): Cycloalkane constituents of the bitumen from Green River Shale. – *Geochimica et Cosmochimica Acta* 35, 661–678.
- CHADEFAUD, M. et EMBERGER, L. (1960): *Traité de Botanique. Systématique. Tome I: Les végétaux non vasculaires. Cryptogamie.* – Masson et Cie Ed., Paris.
- COMBAZ, A. (1980): Les kerogenes vus au microscope. In: *Kerogen insoluble organic matter from sedimentary rocks*, ed.: DURAND, B., 56–112, Edition Technip., Paris.
- DOUGLAS, A. G., EGLINGTON, G. and MAXWELL, J. R. (1969): The hydrocarbons of coorongite. – *Geochimica et Cosmochimica Acta* 33, 569–577.
- FRÉMY, P. et DANGEARD, L. (1938): Observations sur le *Botryococcus braunii* KÜTZING actuel et fossile. – *Ann. Paleont.* 27, 115–136.
- GLIKSON, M., LINDSAY, K. and SAXBY, J. (1989): *Botryococcus* – a planctonic green alga, the source of petroleum through the ages: TEM studies of oil shales and petroleum source rocks. – *Org. Geoch.* 14, 505–608.
- HAJÓS, M. (1976): A pulai Put-3 számú fúrás felső-pannóniai képződményeinek Diatóma flórája. – MÁFI évi jel. az 1974 évről, 263–277.
- HETÉNYI, M. (1985): Organic geochemical features of the maar-type oil shales of Hungary. – *Acta Miner. Petr. Szeged.* 27, 145–152.
- HETÉNYI, M. (1987–1988): Method for measuring the maturity of organic matter in diagenesis stage. – *Acta Miner. Petr. Szeged.* 29, 107–118.
- HETÉNYI, M. and PÁPAY, L. (1986): Type and evolution stage of Hungarian oil shale kerogens. – *Acta Miner. Petr. Szeged.* 28, 109–116.
- JARZEN, D. M. (1978): Some Maestrichtian palynomorphs and their phytogeographical and paleoecological implications. – *Palynology* 2, 29–38.
- JÁMBOR, Á. (1980): A Dunántúli-középhegység pannóniai képződményei. Pulai alginít tagozat. – MÁFI évkönyve 62, 125–129.

- JÁMBOR, Á. and SOLTÍ, G. (1975): Geological conditions of the Upper Pannonian oil-shale deposit recovered in the Balaton Highland and the Kemeneshát. – *Acta Miner. Petr. Szeged.* 27, 9–28.
- KEDVES, M. (1983): Etude paléobotanique sur les schistes pétrolifères du Tertiaire supérieur de Hongrie. – *Revue de Micropaléontologie* 26, 48–53.
- KEDVES, M. (1986a): Dégénération expérimentale des colonies du genre *Botryococcus* des schistes pétrolifères du Tertiaire supérieur de Hongrie. – *Acta Biol. Szeged.* 32, 39–45.
- KEDVES, M. (1986b): Komplex (LM; TEM és vékonyréteg kromatográfiás) vizsgálatok olajpala növényi mikrofossziliáin. – *Bot. Közlem.* 73, 25–32.
- KEDVES, M. (1988): Degradation of the sporoderm under natural and in vitro conditions. – *Acta Biol. Szeged.* 34, 59–69.
- KISS, I. (1939): Die Mikrovegetation der Natrongewässer des Comit. Békés. – *Folia Cryptogamica* 4, 217–266.
- NARAYANA RAO, S. R. and MISRA, S. S. (1949): An oil-bearing alga from the Palana lignite (?Eocene) of Rajputana. – *Curr. Sci.* 18, 380–381.
- NAGY, E. (1976): A dunántúli olajpala-kutató fúrások rétegsorának palinológiai vizsgálata. – *MÁFI évi jel. az 1974. évről*, 247–263.
- POTONIÉ, R. and REHNELT, K. (1971): On the aromatisation of sporin of carboniferous *Lycopside*s. In: Sporopollenin, eds.: BROOKS, J., GRANT, P. R., MUIR, M., VAN GUZEL, R. and SHAW, G., 130–173, Academic Press, London, New York.
- SOLTÍ, G. (1981): A pulai gejzirit. – *MÁFI évi jel. az 1979. évről*, 241–247.
- TRAVERSE, A. (1955): Pollen Analysis of the Brandon Lignite of Vermont. – Bureau of Mines Investigations 5151, U. S. Dept. of the Interior, Washington D. C.
- WILLE, N. (1910): *Botryococcaceae*. In: ENGLER und PRANTL: Die natürlichen Pflanzenfamilien, 241 u. 242 Lief., 32–35, Verlag von WILHELM ENGELMANN, Leipzig.

Appendix

- GUY-OHLSON, D. (1992): *Botryococcus* as an aid in the interpretation of paleoenvironment and depositional processes. – *Rev. Palaeobot. Palynol.* 71, 1–15.

2. DOROG TYPE (MIDDLE EOCENE) SPORE-POLLEN ASSEMBLAGE IN THE SOCKA BEDS OF SLOVENIA I

B. JELEN¹, M. KEDVES², D. SKABERNE¹, A. BREZIGAR¹, S. BUSER³, F. CIMERMAN⁴, K. DROBNE⁴, M. MONOSTORI⁵, R. PAVLOVEC³ and J. PAVŠIČ³

1. Geological Survey of Ljubljana, Institute for Geology, Geotechnics and Geophysics, Dimičeva 14, 61009 Ljubljana, Slovenia, 2. Cell Biological and Evolutionary Micropaleontological Laboratory of the Department of Botany of the J. A. University, H-6701, P. O. Box 657, Szeged, Hungary, 3. Department of Geology and Paleontology, University of Ljubljana, Aškerčeva 2, 61001 Ljubljana, Slovenia, 4. Paleontological Institute Ivan Rakovec, ZRC SAZU, P. O. Box 323, 61001 Ljubljana, Slovenia, 5. Department of Paleontology Eötvös Loránd University, H-1445, P. O. Box 329, Budapest, Hungary

Abstract

During our multidisciplinary studies of the problem of the Oligocene and Lower Miocene stratigraphy in Slovenia, we have established the following: 1. The Socka Beds are heterochronous. 2. The type locality is of Middle Eocene age. 3. The tropical coal forming vegetation was reconstructed on the basis of palynological data. 4. The dominance of the pollen grains associated with spores of tropical ferns (*Anemia*, *Lygodium*, *Gleicheniaceae*, etc.) is identical with the Dorog Basin type. This result changes the previous paleophytogeographical concepts on this subject.

Key words: Palynostratigraphy, Paleophytocology, Paleophytogeography, Middle Eocene, Slovenia.

Introduction

The lithostratigraphic term Socka Beds (Sotzka-Schichten) was introduced by ROLLE in 1858.

The Socka Beds stratigraphic position has never been successfully solved. Namely, stratigraphers found themselves in persistent contradictions as they stated whether at different sites or all embracingly, the Latdorfian (Lower Oligocene), the Rupelian (Middle Oligocene), the Egerian, the Helvetian (Ottangian to Karpatian) age of the Socka Beds, or even vaguely supposed their Eocene age. If we carefully consider the existing literature, it becomes obvious that every work and in particular every rearrangement to accommodate the Socka Beds stratigraphic position to new developments in the chronostratigraphic scale, for example to the establishment of Oligocene or to the proposal of new Paratethian stages, only apparently moved the problem closer to the solution. But, in fact they only added to the confusion concerning the age and the correlation of the Socka Beds in their own way.

The aim of this paper is to present the most important sporomorphs from four localities: Socka, Dobrna, Velenje and Lepena in the Southern Karavanke, and of their comparison with the Transdanubian assemblages in Hungary.

Results

DINOPHYCEAE

Cordosphaeridium cf. *spinosum* var. *deflandrei* (LEJ.-CARPENTIER 1941) DE CON. 1965 (Plate 2.1., figs. 1,2), *C. cf. tiara centrocarpum* (DEFL. and COOKS. 1955) MORG. 1966 (Plate 2.1., figs. 3,4), *Glaphyrocysta* v. *Geridocysta* sp. 1 (Plate 2.1., figs. 5,6), *Glaphyrocysta* v. *Geridocysta* sp.2 (Plate 2.1., figs. 7,8).

PTERIDOPHYTA

LYCOPSIDA

Lycopodium v. *Selaginella*. – *Zlivisporis dorogensis* (KDS. 1965) PACLT. and SICS. 1970 (Plate 2.2., fig. 3).

Selaginellaceae, *Selaginella*. – Cf. *Echinatisporis longechinus* W. KR. 1959 (Plate 2.1., fig. 18).

PTEROPSIDA

Osmundaceae. – *Baculatisporites ovalis* KDS. 1973 (Plate 2.2., fig. 2).

Schizaeaceae

Cf. *Lygodium*. – *Leiotriletes dorogensis* (KDS. 1960) KDS. 1961 fvar. *triplan* (Plate 2.1., fig. 9), *Leiotriletes adriennis* (R. POT. and GELL. 1933) W. KR. 1959 (Plate 2.1., fig. 10), *Leiotriletes microadriennis* W. KR. 1959 (Plate 2.1., fig. 11).

Anemia. – *Cicatricosisporites dorogensis* (R. POT. and GELL. 1933) subfsp. *dorogensis* (Plate 2.1., fig. 9, plate 2.2., figs. 4,5).

Gleicheniaceae

Toroisporis (*Toroisporis*) *teuplitzensis* W. KR. 1962 subfsp. *mediocris* W. KR. 1962 (Plate 2.1., figs. 13,14), *Toroisporis* (*Toroisporis*) *guineti* KDS. 1973 (Plate 2.1., fig. 15), *Concavisporites* (*Concavisporites*) *hungaricus* KDS. 1973 (Plate 2.1., figs. 16,17).

Pteridaceae

Polypodiaceoisporites bauxitus KDS. and J. RÁK. 1965 (Plate 2.2., figs. 6,7), *Polypodiaceoisporites undulosculptatus* KDS. 1973 (Plate 2.2., fig. 8), *Polypodiaceoisporites marxheimensis* (MÜRR. and PF. 1952) W. KR. 1959 (Plate 2.2., figs. 19,20), *Segmentizonosporites palaeogenicus* KDS. and J. RÁK. 1965 (Plate 2.2., figs. 11,12).

Polypodiaceae

Laevigatosporites haardti (R. POT. and VEN. 1934) TH. and PF. 1953 subfsp. *haardti* (Plate 2.2., fig. 15), *Verrucatosporites favus* (R. POT. 1931) TH. and PF. 1953 subfsp. *favus* (Plate 2.2., figs. 16,17).

INCERTAE

Punctatisporites luteticus W. KR. 1959 (Plate 2.1., fig. 12), *Trilites* fsp. (Plate 2.2., figs. 13,14), *Psophosphaera reissingeri* (KDS. 1961) KDS. 1974 (Plate 2.2., figs. 24,25).

GYMNOSPERMATOPHYTA

Cycadopsida, Ginkgoaceae v. Cycadaceae. – *Cycadopites kyushuensis* (TAKAHASHI 1961) KDS. 1968 (Plate 2.3., figs. 1,2)

Coniferophytina

?Abietaceae, *Pseudotsuga* v. *Larix*. – Cf. *Psophosphaera* fsp. (Plate 2.2., figs. 22,23).

Taxodiaceae v. Cupressaceae. – *Inaperturopollenites concedipites* (WODEH. 1933) W. KR. 1971 (Plate 2.2., fig. 1).

Araucariaceae

Araucariacites balinkaense KDS. 1974 (Plate 2.2., figs. 20,21).

? Coniferophytina

Cupressacites dorogensis (KDS. 1961) KDS. 1974 (Plate 2.2., figs. 26,27).

Reworked *gymnosperm* pollen grain. – *Alisporites* fsp. (Plate 2.2., figs. 18,19).

ANGIOSPERMATOPHYTA

DICOTYLEDONOPSIDA

Fagaceae. – *Fususpollenites fusus* (R. POT. 1934) KDS. 1978, *Fagaceae* (Plate 2.3., figs. 19,20).

Myricaceae. – *Plicapollis pseudoexcelsus* (W. KR. 1958) W. KR. 1961 subfsp. *turgidus* PF. 1953 (Plate 2.3., figs. 27,28).

Juglandaceae. – Cf. *Platycaryapollenites* fsp. (Plate 2.3., figs. 29,30), *Plicatopollis lunatus* KDS. 1974 (Plate 2.3., figs. 31,32), *Plicatopollis hungaricus* KDS. 1974 (Plate 2.3., figs. 33,34), *Plicatopollis krutzschii* KDS. 1974 (Plate 2.3., figs. 35,36).

Icacinaceae. – *Compositoipollenites rhizophorus* (R. POT. 1934) R. POT. 1960 subfsp. *rhizophorus* (Plate 2.3., figs. 39,40).

Elaeagnaceae v. Simarubaceae. – *Pentapollenites laevigatus* W. KR. 1962a subfsp. *laevigatoides* W. KR. 1962a (Plate 2.3., figs. 25,26).

Aquifoliaceae. – Cf. *Ilexpollenites erdmani* KDS. 1978 (Plate 2.3., figs. 15,16).

Bombacaceae. – *Bombacacidites* fsp. (Plate 2.3., figs. 37,38).

Sapotaceae. – *Tetracolporopollenites urkuticus* KDS. 1978 (Plate 2.3., figs. 21,22).

Calyceraceae. – *Intragranulitricolporites tumescens* (KDS. 1964) KDS. 1978 (Plate 2.3., figs. 17,18).

Incertae. – *Eocaenipollis* fsp. (Plate 2.3., figs. 23,24).

MONOCOTYLEDONOPSIDA

Palmae. – *Monocolpopollenites tranquillus* (R. POT. 1934) TH. and PF. 1953 subfsp. *tranquillus* (Plate 2.3., figs. 3–6).

Nipa. – *Spinizonocolpites* cf. *prominatus* (McINTYRE 1965) KDS. 1974 (Plate 2.3., figs. 7,8), cf. *Nipa*. – *Spinizonocolpites* fsp. (Plate 2.3., figs. 9–12).

North African Upper Cretaceous reworking: *Proxapertites africanus* KDS. nom. nud. (Plate 2.3., figs. 13,14).

Conclusions

1. The presented spore-pollen assemblage is identical with the Middle Eocene “Dorog type” of Hungary.

2. So far the Middle Eocene age is restricted to the coal bearing strata at Socka, Dobrna, Velenje and Lepena (near Jesenice) at Slovenia–Austria–Italy border. All these localities belong to the Southern Karavanka Mountains.

Acknowledgements

This work was supported by the grants as follows. Slovenia: 2 – 2238–215, Hungary: OTKA 1/5, T–007206.

References

- GRUAS-CAVAGNETTO, C. (1968): Étude palynologique des divers gisements du Sparnacien du Bassin de Paris. – Mém. Soc. Géol. de France *110*, 1–144.
- GRUAS-CAVAGNETTO, C. (1976): Les marqueurs stratigraphiques (*Dinoflagellés*) de l'Eocène du Bassin de Paris et de La Manche Orientale. – Rev. de Micropaléont. *18*, 221–228.
- KEDVES, M. (1960): Études palynologiques dans le Bassin de Dorog – I –. – Pollen et Spores *2*, 101–118.
- KEDVES, M. (1961): Études palynologiques dans le Bassin de Dorog – II –. – Pollen et Spores *3*, 101–153.
- KEDVES, M. (1964): Sporomorphes nouveaux des couches éocènes de Hongrie. – Pollen et Spores *6*, 196–201.
- KEDVES, M. (1965): Ein neuer reticulater Sporen-Typ aus dem Eozän des Braunkohlenbeckens von Dorog (Ungarn). – Paläont. Z. *39*, 188–192.
- KEDVES, M. (1968): Études palynologiques des couches du Tertiaire inférieur de la Région Parisienne. III. Pollens inaperturés, à ballonnets, polypliqués, monocarpés, disulqués, trichotomosulqués et proxaperturés. – Pollen et Spores *10*, 315–334.
- KEDVES, M. (1973): Paleogene fossil sporomorphs of the Bakony Mountains Part I. – Stud. Biol. Acad. Sci. Hung. *12*, 1–134.
- KEDVES, M. (1974): Paleogene fossil sporomorphs of the Bakony Mountains Part II. – Stud. Biol. Acad. Sci. Hung. *13*, 1–124.
- KEDVES, M. (1978): Paleogene fossil sporomorphs of the Bakony Mountains Part III. – Stud. Biol. Acad. Sci. Hung. *15*, 1–166.
- KEDVES, M. and RÁKOSY, J. (1965): Zonotrilete microspores from the Eocene bauxite layers Gánt in Hungary. – Acta Biol. Szeged. *11*, 233–244.
- KRUTZSCH, W. (1958): Sporen- und Pollen-Gruppen aus der Oberkreide und dem Tertiär Mitteleuropas und ihre stratigraphische Verteilung. – Z. angew. Geol. *3*, 509–548.
- KRUTZSCH, W. (1959): Mikropaläontologische (sporenpaläontologische) Untersuchungen in der Braunkohle des Geiseltales, I. – Geologie, Beih. *21/22*, 1–425.
- KRUTZSCH, W. (1961): Beitrag zur Sporenpaläontologie der präoberoligozänen kontinentalen und marinen Tertiärlagerungen Brandenburgs. – Berichte der Geol. Ges. *4*, 290–343.
- KRUTZSCH, W. (1962a): Mikropaläontologische (sporenpaläontologische) Untersuchungen in der Braunkohle des Geiseltales. II. Die Formspezies der Pollengattung *Penapollenites* KRUTZSCH 1958. – Paläontologische Abhandlungen *1/2*, 75–103.
- KRUTZSCH, W. (1962b): Atlas der mittel- und jungtertiären dispersen Sporen- und Pollen- sowie der Mikroplanktonformen des nördlichen Mitteleuropas. I. Laevigate und toriate trilete Sporenformen. – VEB Deutscher Verlag der Wissenschaften, Berlin.
- KRUTZSCH, W. (1971): Atlas der mittel- und jungtertiären dispersen Sporen- und Pollen- sowie der Mikroplanktonformen des nördlichen Mitteleuropas VI. *Coniferenpollen* (*Saccites* und "*Inaperturates*"). VEB Gustav Fischer Verlag, Jena.
- MCINTYRE, D. J. (1965): Some new pollen species from New Zealand Tertiary deposits. – N. Z. J. Bot. *3*, 204–214.
- PACLTOVÁ, B. and SIMONCSICS, P. (1970): New types of spores (Genera and species) from the Bohemian Miocene. – Paläont. Abh. B, *3*, 599–617.
- POTONIÉ, R. (1931): Zur Mikroskopie der Braunkohlen. Tertiäre Sporen- und Blütenstaubformen. 4. Mitt. – Z. Braunkohle *30*, H. 27, 554–556.
- POTONIÉ, R. (1934): Zur Mikrobotanik des eozänen Humodils des Geiseltales. – Arb. aus Inst. Paläobotanik u. Petrogr. Brenngesteine *4*, 25–125.
- POTONIÉ, R. (1960): Synopsis der Gattungen der Sporae dispersae, III: Nachträge Sporites, Fortsetzung Pollenites Mit Generalregister zu Teil I–III. – Beih. Geol. Jb. *39*, 1–189.
- POTONIÉ, R. und GELLETICH, J. (1933): Ueber Pteridophytensporen einer eozänen Braunkohle aus Dorog in Ungarn. – Sitz. – Ber. naturf. Fr., 317–328.
- POTONIÉ, R. und VENITZ, H. (1934): Zur Mikrobotanik des miozänen Humodils der niederrheinischen Bucht. – Arb. aus Inst. Paläobotanik u. Petrogr. Brenngesteine *5*, 5–53.

- ROLLE, F. (1958): Über die geologische Stellung der Sotzka-Schichten. – Sitzungber. Akad. Wiss. Math.-naturwiss. Cl. 30, 3–34.
- TAKAHASHI, K. (1961): Pollen und Sporen des westjapanischen Alttertiärs und Miozäns. Teil II. – Mem. Fac. Sci. Kyushu Univ., der. D (Geol.) 11, 279–345.
- THOMSON, P. W. und PRUG, H. D. (1953): Pollen und Sporen des mitteleuropäischen Tertiärs. – Palaeontographica B, 94, 1–139.
- WILSON, G. J. (1988): Paleocene and Eocene Dinoflagellatae Cysts from Waipawa, Hawkes, Bay, New Zealand. – New Zealand Geological Survey Palaeontological Bulletin 57, 1–96.
- WODEHOUSE, R. P. (1933): Tertiary pollen. II. The oil shales of the Eocene Green River Formation. – Bull. Torrey Bot. Club 60, 479–524.

Plate 2.1.

- 1,2. *Cordosphaeridium* cf. *spinosum* var. *deflandrei* (LEJ.-CARPENTIER 1941) DE CON. 1965, slide: Lepena-6-1, cross-table number: 11.9/144.6.
- 3,4. *Cordosphaeridium* cf. *tiara centrocarpum* (DEFL. and COOKS. 1955) MORG. 1966, slide: Lepena-6-1, cross-table number: 9.2/145.8.
- 5,6. *Glaphyrocysta* v. *Geridocysta* sp.1, slide: Lepena-6-1, cross-table number: 9.1/140.1.
- 7,8. *Glaphyrocysta* v. *Geridocysta* sp.2, slide: Lepena-6-1, cross-table number: 19.9/139.9.
9. *Leiotriletes dorogensis* (KDS. 1960) KDS. 1961 fvar. *triplan*, *Schizaeaceae*, cf. *Lygodium*, and *Cicatricosisporites dorogensis* R. POT. and GELL. 1933. subfsp. *dorogensis*, *Schizaeaceae*, *Anemia*; slide: Velenje Sv. Bric-3-1, cross table number: 5.4/133.5.
10. *Leiotriletes adriennis* (R. POT. and GELL. 1933) W. KR. 1959, *Schizaeaceae* cf. *Lygodium*; slide: Dobrna-2-1, 9.3/130.9.
11. *Leiotriletes microadriennis* W. KR. 1959, *Schizaeaceae* cf. *Lygodium*; slide: Dobrna-2-1, cross-table number: 5.6/144.1.
12. *Punctatisporites luteticus* W. KR. 1959; slide: Velenje Sv. Bric-4-2, cross-table number: 6.9/137.1.
- 13,14. *Toroisporis* (*Toroisporis*) *teuplitzensis* W. KR. 1962 subfsp. *mediocris* W. KR. 1962. cf. *Gleicheniaceae*, slide: Dobrna-2-1, cross-table number: 23.2/147.2.
15. *Toroisporis* (*Toroisporis*) *guineti* KDS. 1973, cf. *Gleicheniaceae*, slide: Velenje Sv. Bric-3-1, cross-table number: 18.2/132.4.
- 16,17. *Concavisporites* (*Concavisporites*) *hungaricus* KDS. 1973, *Gleicheniaceae*, slide: Velenje Sv. Bric-4-2, cross-table number: 17.2/135.2.
18. Cf. *Echinatisporis longechinus* W. KR. 1959, *Selaginellaceae*, *Selaginella*, slide: Lepena-3-1, cross-table number: 13.5/135.6.

Plate 2.2.

1. *Punctatisporites luteticus* W. KR. 1959, with *Inaperturopollenites concedipites* (WODEH. 1933) W. KR. 1971, *Taxodiaceae-Cupressaceae*, slide: Velenje Sv. Bric-6-1, cross-table number: 11.4/129.6.
2. *Baculatisporites ovalis* KDS. 1973, *Osmundaceae*, slide: Dobrna-2-1, cross-table number: 24.8/134.3.
3. *Zlivisporis dorogensis* (KDS. 1965) PACLT. and SICS. 1970, *Lycopodiaceae*, *Lycopodium* v. *Selaginella*, Velenje Sv. Bric-4-2, cross-table number 22.3/129.6.
- 4,5. *Cicatricosisporites dorogensis* R. POT. and GELL. 1933 subfsp. *dorogensis*, *Schizaeaceae*, *Anemia*, slide: Dobrna-2-1, cross-table number: 10.2/142.3.
- 6,7. *Polypodiaceoisporites bauxianus* KDS. and J. RÁK. 1965, slide: Lepena-3-1, cross-table number: 20.8/146.7.
8. *Polypodiaceoisporites undulosculptatus* KDS 1973, *Pteridaceae*, slide: Velenje Sv. Bric-4-2, cross-table number: 6.5/147.9.
- 9,10. *Polypodiaceoisporites marxheimensis* (MÜRR. and PF. 1952) W. KR. 1959, *Pteridaceae*, slide: Velenje Sv. Bric-6-1, cross-table number: 9.5/137.2.

- 11,12. *Segmentizonosporites palaeogenicus* KDS. and J. RÁK. 1965, slide: Velenje Sv. Bric-4-1, cross-table number: 11.3/140.8.
- 13,14. *Trilites* fsp., slide: Velenje Sv. Bric-4-2, cross-table number: 12.7/146.2.
15. *Laevigatosporites haardii* (R. POT. and VEN. 1934) TH. and PF. 1953 subfsp. *haardii*, *Polypodiaceae*, slide: Dobrna-1-1, cross-table number: 8.6/132.1.
- 16,17. *Verrucatosporites favus* (R. POT. 1931) TH. and PF. 1953 subfsp. *favus*, *Polypodiaceae*, slide: Velenje Sv. Bric-6-1, cross-table number: 9.5/129.6.
- 18,19. *Alisporites* fsp. – reworked, slide: Socka-1-1, cross-table number: 7.7/144.8.
- 20,21. *Araucariacites balinkaense* KDS. 1974, *Araucariaceae*, slide: Dobrna-1-1, cross-table number: 8.9/130.8.
- 22,23. Cf. *Psophosphaera* fsp., ?*Abietaceae*, *Pseudotsuga* v. *Larix*, slide: Dobrna-1-1, cross-table number: 4.8/140.3.
- 24,25. *Psophosphaera reissingeri* (KDS. 1961) KDS. 1974, slide: Velenje Sv. Bric-6-1, cross-table number: 5.2/142.6.
- 26,27. *Cupressacites dorogensis* (KDS. 1961) KDS. 1974, cf. *Coniferae*, slide: Lepena-8-1, cross-table number: 7.8/141.8.

Plate 2.3.

- 1,2. *Cycadopites kyushuensis* (TAKAHASHI 1961) KDS. 1968, *Ginkgoaceae*, v. *Cycadaceae*, slide: Velenje Sv. Bric-3-1, cross-table number: 7.8/130.6.
- 3,4. *Monocolpopollenites tranquillus* (R. POT. 1934) TH. and PF. 1953. subfsp. *tranquillus*, *Palmae*, slide: Velenje Sv. Bric-6-1, cross-table number: 17.2/151.3.
- 5,6. *Monocolpopollenites tranquillus* (R. POT. 1934) TH. PF. 1953 subfsp. *tranquillus*, *Palmae*, slide: Velenje Sv. Bric-6-1, cross-table number: 5.2/151.2.
- 7,8. *Spinizonocolpites* cf. *prominatus* (MCINTYRE 1965) KDS. 1974, *Palmae*, *Nipa*, slide: Lepena-3-1, cross-table number: 9.2/145.2.
- 9,10. *Spinizonocolpites* fsp., *Palmae*, cf. *Nipa*, slide: Lepena-2-1, cross-table number: 17.2/143.2.
- 11,12. *Spinizonocolpites* fsp., *Palmae*, cf. *Nipa*, slide: Lepena-3-1, cross-table number: 5.8/144.2.
- 13,14. "*Proxapertites africanus* KDS. nom. nud.", slide: Velenje Sv. Bric-3-1, cross-table number: 5.9/131.9.
- 15,16. Cf. *Ilexpollenites erdtmani* KDS. 1978, *Aquifoliaceae*, slide: Velenje Sv. Bric-3-1, cross-table number: 15.2/142.3.
- 17,18. *Intragranulitricolporites tumescens* (KDS. 1964) KDS. 1978, *Calyceraceae*, slide: Dobrna-1-1, cross-table number: 10.1/135.8.
- 19,20. *Fususpollenites fusus* (R. POT. 1934) KDS. 1978, *Fagaceae*, *Castanopsis*, slide: Dobrna-1-1, cross-table number: 8.9/139.8.
- 21,22. *Tetracolporopollenites urkauticus* KDS. 1978, *Sapotaceae*, slide: Dobrna-1-1, cross-table number: 8.1/140.1.
- 23,24. *Eocaenipollis* fsp., slide: Dobrna-1-1, cross-table number: 15.6/135.1.
- 25,26. *Pentapollenites laevigatus* W. KR. 1962 subfsp. *laevigatoides* W. KR. 1962, *Elaeagnaceae* v. *Simarubaceae*, slide: Velenje Sv. Bric-3-1, cross-table number: 16.3/144.4.
- 27,28. *Plicatopollis pseudoexcelsus* (W. KR. 1958) W. KR. 1961 subfsp. *nurgidus* PF. 1953, cf. *Myricaceae*, slide: Dobrna-1-1, cross-table number: 8.9/127.6.
- 29,30. Cf. *Platycaryapollenites* fsp., *Juglandaceae*, slide: Velenje Sv. Bric-3-1, cross-table number: 7.2/135.9.
- 31,32. *Plicatopollis lunatus* KDS. 1974, *Juglandaceae*, slide: Velenje Sv. Bric-3-1, cross-table number: 8.6/130.1.
- 33,34. *Plicatopollis hungaricus* KDS. 1974, *Juglandaceae*, slide: Dobrna-1-1, cross-table number: 8.9/131.2.
- 35,36. *Plicatopollis krutzschii* KDS. 1974, *Juglandaceae*, slide: Velenje Sv. Bric-3-1, cross-table number: 9.3/143.8.
- 37,38. *Bombacacidites* fsp., *Bombacaceae*, slide: Lepena-3-1, cross-table number: 10.6/136.1.
- 39,40. *Compositoipollenites rhizophorus* (R. POT. 1934) R. POT. 1960 subfsp. *rhizophorus*, *Icacinaceae*, slide: Lepena-8-1, cross-table number: 10.2/133.1.




Plate 2.1.




Plate 2.2.




Plate 2.3.

3. PLANT MICROFOSSILS FROM THE UPPER CRETACEOUS AND LOWER TERTIARY LAYERS OF NORTHERN SPAIN I.

M. KEDVES

Cell Biological and Evolutionary Micropaleontological Laboratory of the Department of Botany of the J. A. University, H-6701, P. O. Box 657, Szeged, Hungary

Abstract

The spore-pollen assemblages of two localities of the Upper Cretaceous layers from Northern Spain are presented in this contribution. In the sample of the Zambrana section (30 km South of Miranda de Ebro) the *gymnospermous Classoidites glandis* occurs in a dominant quantity with a large number of early *Brevaxonate* pollen grains: *Atlantopollis* and *Complexiopollis* fssp. This is similar to the spore-pollen assemblage isolated from Oviedo (SOLÉ DE PORTA, 1978) and to those of Fermetelos, Oia, etc. from Portugal (KEDVES and PÁRDUTZ, 1983). As for their geological age, the Lower Turonian seems to be probable. The spore-pollen assemblage of the Sobron section (80 km North-West of Burgos) is quite different. Dominant forms are the disaccate *gymnosperm* pollen grains. The quantity of the evolved early *Brevaxonate* pollen grains (*Eunormapollis*: *Papillopollis*, *Interporopollenites*, *Verruculopollis*, *Endopollis*, *Krutzschipollis*, etc.) and that of the *Postnormapollis* is relatively large. A similar palynomorph assemblage was described from Guadalix (MÉDUS and ALVAREZ RAMIS, 1989), its geological age is probably Santonian.

Key words: Palynology, fossil, Upper Cretaceous, Spain.


Introduction

There are several publications concerning the spore-pollen assemblages of the Upper Cretaceous sediments of Spain. On the basis of the different concepts of these researches the following, previously published papers may be mentioned:

1. Description of basic plant microfossil remains. AMEROM (1965, Cenomanian-Turonian of the Province of León, Northern Spain), SOLÉ DE PORTA (1978, Cenomanian from Los Alrededores de Oviedo), ALVAREZ RAMIS (1985, El Vellón, Madrid), PORTA, KEDVES, SOLÉ DE PORTA and CIVIS (1985, Maastrichtian of Barranco de La Posa, Lérida), ALVAREZ RAMIS, CLEMENTE BELMONTE, FERNÁNDEZ MARRÓN, GALÁN RUIZ and GÓMEZ PORTER (1987, El Vellón, Madrid), MÉDUS (1987, Campanian from Cedano, N. of Burgos), MÉDUS and ALVAREZ RAMIS (1989, Senonian from Guadalix of the Province of Madrid), ALVAREZ RAMIS and DOUBINGER (1980, Torrelaguna, Madrid).
2. Comparative investigations; Northern Spain with the Southern part of France; MÉDUS (1970a,b, 1972, 1975, 1978).

3. Paleophytogeography concerning the Mediterranean Region including the Iberian Peninsula: KRUTZSCH (1967, in: GÓCZÁN, GROOT, KRUTZSCH and PACLTOVÁ), KEDVES and DINIZ (1983), KEDVES (1985).

4. Investigation of the in situ pollen grains from Upper Cretaceous sediments of Spain; ALVAREZ RAMIS (1980, 1981), ALVAREZ RAMIS and FERNÁNDEZ MARRÓN (1986), ALVAREZ RAMIS, BARALE and FERNÁNDEZ MARRÓN (1987) BARALE, FERNÁNDEZ MARRÓN and ALVAREZ RAMIS (1988).



Text-fig. 3.1.

Schematical map of the sampling localities following MARTINS.

Materials and Methods

Dr. ULRICH P. MARTINS (Department of Geology and Paleontology of the University of Tübingen, R. F. Germany) have sent me slides for palynological investigations. The materials of investigations may be divided into two groups. This paper deals with the spore-pollen assemblages of the lower two levels. On the basis of the letter of Dr. U. P. MARTINS dated on 3. 5. 1987, the data of the marlstone samples are the following.

Sample 194/10 (2 slides) came from the Sobron Section, 80 km North-West of Burgos, approximately Lower Santonian in age.

Sample 268/14 (4 slides) originates from the Zambrana Section, 30 km South of Miranda de Ebro, same age. The treatment of the samples started with HNO₃ (65%).

The localities of the sampling by Dr. U. P. MARTINS are illustrated on text-fig. 3.1. (cf. MARTINS 1988, 1989). The stratigraphic column of Quintanilla la Ojada – Losa Valley – (Text-fig. 3.2.) illustrates the stratigraphic situations of the samples under palynological investigation. On the basis of the preliminary studies, the two levels presented in this contribution represents the oldest part of the Upper Cretaceous in this region. For the first step the spore-pollen assemblages are elaborated, the *Acritarchs* will be the subjects of further investigations.

Results

1. ZAMBRANA SECTION (30 KM SOUTH OF MIRANDA DE EBRO)

No: 268/14

Quantitative data


	No of specimens
<i>Hystriospheraeidae</i>	196
<i>Sporites</i>	7
<i>Disaccites</i>	8
<i>Classoidites</i>	57
<i>Atlantopollis</i>	43
<i>Complexiopollis</i>	17

Qualitative data

Sporites

Stereisporites aegypticus KEDVES 1986, *Sphagnaceae* (Plate 3.1., figs. 1,2),
Ischiosporites cf. *tuberosus* DÖRING 1965 (Plate 3.1., figs. 10,11),
Ischiosporites cf. *pseudoreticulatus* (COUPER 1958) DÖRING 1965 (Plate 3.1., fig. 12),
Cicatricosisporites baconicus H. DEÁK 1963, *Schizaeaceae* (Plate 3.1., fig. 3),
Appendicisporites potomacensis BRENNER 1963, *Schizaeaceae* (Plate 3.1., fig. 4),
Cf. *Clavatisporites* fsp. (Plate 3.1., fig. 5),

QUINTANILLA LA OJADA - LOSA VALLEY



◀ Text-fig. 3.2.

Stratigraphical column of Quintanilla La Ojada - Losa Valley and the position of the samples which are the subjects of this research program. Schema following MARTINS.

Liburnisporites fsp. (Plate 3.1., figs. 6,7),
Uvaesporites cf. *neerlandicus* HERNGREEN, FELDER, KEDVES and MEESSEN 1986 (Plate 3.1., fig. 13),
Camarozonosporites insignis NORRIS 1967, *Lycopodiaceae* (Plate 3.1., figs. 8,9).

Pollenites

Podocarpidites multesimus (BOLKHOVITINA 1956) POCKOCK 1962 (Plate 3.1., figs. 14,15),
Classoidites glandis AMEROM 1965 (Plate 3.1., figs. 16,17),
Atlantopollis microreticulatus KRUTZSCH 1967 (Plate 3.1., figs. 18,19),
Atlantopollis vilaflorensis KEDVES and DINIZ 1979 (Plate 3.1., figs. 20,21),
Atlantopollis grooti KEDVES and DINIZ 1979 (Plate 3.1., figs. 22,23),
Atlantopollis choffatii DINIZ, KEDVES and SIMONCSICS 1974 (Plate 3.1., fig. 24),
Complexiopollis praeatumescentes KRUTZSCH 1959 (Plate 3.1., figs. 25–28),
Complexiopollis vulgaris (GROOT and GROOT 1962) GROOT and KRUTZSCH 1967 (Plate 3.1., figs. 29,30),
Complexiopollis subtilis (KRUTZSCH 1959) KRUTZSCH 1967 (Plate 3.1., figs. 31,32),
Complexiopollis latis (KRUTZSCH 1959) KRUTZSCH 1967 (Plate 3.1., figs. 33,34).

2. SOBRON SECTION (80 KM. NORTH-WEST OF BURGOS)

No: 194/10

Quantitative data

	No of specimens
<i>Hystriospheraeidae</i>	124
<i>Dissaccites</i>	89
<i>Classoidites</i>	2
<i>Longaxones</i>	3
<i>Eunormapolles</i>	52
<i>Postnormapolles</i>	6

Qualitative data

Pollenites

Pinuspollenites spherisaccus BRENNER 1963 (Plate 3.2., figs. 1,2),
Podocarpidites verruculatus (TREVISAN 1967) KRUTZSCH 1971 (Plate 3.2., figs. 3,4),
Retitricolpites minutus (BRENNER 1963) DETTMANN 1973 (Plate 3.2., figs. 5,6),
Illexpollenites erdtmani KEDVES 1978, *Aquifoliaceae* (Plate 3.2., figs. 7,8),
Complexiopollis cf. *patulus* TSCHUDY 1973 (Plate 3.3., figs. 31,32),
Verruculopollis fsp. (Plate 3.2., figs. 9,10),
Endopollis fsp. (Plate 3.2., figs. 11,12),
Krutzschipollis spatiosus GÓCZÁN (Plate 3.2., figs. 13,14),

Semiculopollis medius KEDVES and DINIZ 1983 (Plate 3.2., figs. 15,16),
Papillopollis fsp. (Plate 3.2., figs. 17,18),
Papillopollis pflugii KEDVES and PITTAU 1979 (Plate 3.3., figs. 1,2),
Papillopollis vancampoae KEDVES and PITTAU 1979 (Plate 3.3., figs. 3–6),
Papillopollis rugulatus KEDVES and PITTAU 1979 (Plate 3.3., figs. 7,8),
Papillopollis regulus PFLUG 1953 (Plate 3.3., figs. 9,10),
Rocheipollenites triangulus KEDVES and DINIZ 1980 (Plate 3.3., figs. 11,12),
Triangulipollis cf. *magnus* KEDVES and DINIZ 1980–1981 (Plate 3.3., figs. 15,16),
Interporopollenites plicatus KEDVES and HEGEDÜS 1975 (Plate 3.3., figs. 13,14,17,18),
Interporopollenites stanley KEDVES and HEGEDÜS 1975 (Plate 3.3., figs. 19,20),
Interporopollenites concavus KEDVES and HEGEDÜS 1975 (Plate 3.3., figs. 21,22),
Interporopollenites dinizae KEDVES and HEGEDÜS 1975 (Plate 3.3., figs. 23,24),
Interporopollenites ornatus KEDVES and HEGEDÜS 1975 (Plate 3.3., figs. 25,26),
Alnipollenites fsp. (Plate 3.3., figs. 27,28),
Triporopollenites nointelensis KEDVES 1970, *Corylaceae*, (Plate 3.3., figs. 29,30),
Triatriopollenites lubomirovae (GLADKOVA 1965) KEDVES 1974, *Myricaceae* (Plate 3.3., figs. 31,32),
Inaperturopollenites lacunosus VAN HOEKEN-KLINKENBERG 1964 (Plate 3.3., figs. 33,34).

Discussion and Conclusions

1. The large quantity of the planctonic organisms in both assemblages need further investigations in the future.
2. The spore-pollen assemblage of the Zambrana section is a very good basis of comparison for the lower part of the Upper Cretaceous spore-pollen assemblages in Northern Spain. This can be characterized with the early *Normapolles* genera, *Atlantopollis* and *Complexiopollis*. The lack of the *Probrevaxones*, and the fact, that during my investigation *Longaxones* pollen grains were not observed is worth mentioning. Obviously, further investigations will probably enrich our knowledge of this matter. This assemblage may be compared to further European localities in the boreal region, too; cf. KRUTZSCH 1967, in GÓCZÁN, GROOT, KRUTZSCH and PACLTOVÁ.
3. The spore-pollen assemblage of the Sobron locality is much younger, with no intermediate assemblages till this time. This is similar to the Upper Cretaceous spore-pollen assemblages from Portugal. Until this time we have been able to use the so-called Aveiro type from Portugal for comparison, but further investigations are necessary needed.

Acknowledgements

This work was supported by the Grant OTKA 1/5, T 007206.

References

- ALVAREZ-RAMIS, C. (1980): *Frenelopsis* from the Upper Cretaceous of Torrelaguna, Spain. – Int. Paleobot. Conf. Reading, Abstracts, 1.
- ALVAREZ-RAMIS, C. (1981): Estudio de *Frenelopsis oligostomata* (ROMARIZ) ALVIN, procedente del Cretácico superior de Torrelaguna (Madrid). – Estudios geol. 37, 83–87.
- ALVAREZ-RAMIS, C. (1985): Sur la paléocologie des alentours du barrage "El Vellon" (Madrid, Espagne) pendant la Crétacé supérieur. – 110^e Congr. nat. des Soc. savantes Montpellier, sciences 5, 115–125.
- ALVAREZ-RAMIS, C., BARALE, G. and FERNÁNDEZ MARRÓN, T. (1987): Cones containing pollen of *Classopollis* type of *Frenelopsis oligostomata* ROMARIZ emend. ALVIN 1977 (*Cheirolepidaceae*) from Upper Cretaceous of Madrid Province. – XIV. Int. Bot. Congr. Berlin (West), Germany, Abstracts, 332.
- ALVAREZ RAMIS, C., CLEMENTE BELMONTE, P., FERNÁNDEZ MARRÓN, T., GALÁN RUIZ, C. y GÓMEZ PORTER, P. (1987): Formas de *Normapolles* del Senoniense del Embalse de "El Vellon" Madrid. – VII Bial de la Real Soc. Española de Hist. Nat., Pamplona, Resúmenes, 130.
- ALVAREZ RAMIS, C. y DOUBINGER, J. (1980): Los microfósiles vegetales del Cretácico superior de Torrelaguna (Madrid). – Bol. R. Soc. Española Hist. Nat. (Geol.) 78, 205–215.
- ALVAREZ RAMIS, C. y FERNÁNDEZ MARRÓN, T. (1986): Identification de *Classopollis classoides* (PFLUG) POCKOCK y JANSONIUS 1961 hallados en el supracretácico de la Provincia de Madrid. – VI Simposio de Palinología, Salamanca, Resúmenes, 84.
- AMEROM VAN, H. W. J. (1965): Upper Cretaceous pollen and spores assemblages from the so-called "Wealden" of the Province of Leon (Northern Spain). – Pollen et Spores 7, 93–133.
- BARALE, G., FERNÁNDEZ MARRÓN, T. et ALVAREZ-RAMIS, C. (1988): Étude des cônes mâles de *Frenelopsis* SCHENK. emend. REYMANOWNA & WATSON, 1976 (*Cheirolepidaceae*) du Crétacé supérieur de Torrelaguna (Province de Madrid – Espagne). – Geobios 21, 187–199.
- BOLKHOVITINA, N. A. (1956): Atlas of Spores and Pollen from Jurassic and Lower Cretaceous Deposits of the Vilyui Depression (Russian). – Trans. Geol. Inst. Acad. Sci. U.S.S.R., 2, 1–118.
- BRENNER, G. J. (1963): The spores and pollen of the Potomac Group of Maryland. – Dept. of Geol. Min. and Water Res. Bull. 27, 1–215.
- COUPER, R. A. (1958): British Mesozoic Microspore and Pollen Grains, A Systematic and Stratigraphic Study. – Palaeontographica B, 103, 75–179.
- DEÁK, H. M. (1963): Quelques spores striées de l'étage Aptien. – Rev. de Micropaléontologie 5, 251–256.
- DETTMANN, M. E. (1973): Angiospermous pollen from Albian to Turonian sediments of eastern Australia. – Geol. Soc. of Australia, Special Publication 4, 3–34.
- DINIZ, F., KEDVES, M. et SIMONCSICS, P. (1977): Les sporomorphes principaux de sédiments crétacés de Vila Flor et de Carrajão, Portugal. – Com. dos Serv. Geol. de Portugal 58, 161–178.
- DÖRING, H. (1965): Die sporenpaläontologische Gliederung des Wealden in Westmecklenburg (Struktur Werle). – Geologie 14, BH 47, 1–118.
- GLADKOVA, A. N. (1965): Fossil *Myricaceae* of Western Siberia (Russian). – Tr. VNIGRI 239, 142–190.
- GÓCZÁN, F., GROOT, J. J., KRUTZSCH, W. und PACLTÓVÁ, B. (1967): Die Gattungen des "Stemma *Normapolles* PFLUG 1953 b" (*Angiospermae*) Neubeschreibungen und Revision europäischer Formen (Oberkreide bis Eozän). – Paläont. Abh. 2, 427–633.
- GÓCZÁN, F. (1967): In GÓCZÁN, F., GROOT, J. J., KRUTZSCH, W. und PACLTÓVÁ, B.
- GROOT, J. J. and GROOT, C. R. (1962): Plant Microfossils from Aptian, Albian and Cenomanian deposits of Portugal. – Com. dos Serv. Geol. de Portugal 46, 133–171.
- GROOT, J. J. and KRUTZSCH, W. (1967): In GÓCZÁN, F., GROOT, J. J., KRUTZSCH, W. und PACLTÓVÁ, B.
- HERNGREEN, G. F. W., FELDER, W. M., KEDVES, M. and MEESSEN, J. P. M. T. (1986): Micropaleontology of the Maestrichtian in Borehole Bunde, The Netherlands. – Rev. Palaeobot. Palynol. 48, 1–70.
- KEDVES, M. (1970): Études palynologiques des couches du Tertiaire inférieur de la Région Parisienne. V. Pollens triporés, subtriporés et intritriporés. – Pollen et Spores 12, 83–97.
- KEDVES, M. (1974): Paleogene fossil sporomorphs of the Bakony Mountains. Part II. – Studia Biol. Acad. Sci. Hung. 13, 1–124.

- KEDVES, M. (1978): Paleogene fossil sporomorphs of the Bakony Mountains. Part III. – *Studia Biol. Acad. Sci. Hung.* 15, 1–166.
- KEDVES, M. (1985): The present day state of Upper Cretaceous palaeophytogeography on palynological evidence. – *Acta Biol. Szeged.* 31, 115–127.
- KEDVES, M. (1986): Palynological investigations of Prequaternary sediments of Egypt. Lower part of the Nubian Sandstone in the Kharga Oasis. – *Z. geol. Wiss. Berlin* 14, 331–355.
- KEDVES, M. et DINIZ, F. (1979): Les pollens d'Angiospermes du Crétacé de Vila Flor, Portugal, Genres de forme *Atlantopollis* et *Limaipollenites*. – *Bol. Soc. Geol. de Portugal* 21, 203–216.
- KEDVES, M. et DINIZ, F. (1980–1981): Contribution à la connaissance des pollens d'Angiospermes du Crétacé supérieur du Portugal. – *Bol. Soc. Geol. de Portugal* 22, 19–39.
- KEDVES, M. et DINIZ, F. (1983): Les *Normapolles* du Crétacé supérieur en Europe: Implications Paléobiogéographiques. – *Geobios* 16, 329–345.
- KEDVES, M. and HEGEDÜS, M. (1975): Pollen grains of the *Interporopollenites* fgen. from sediments of the Upper Cretaceous period in Portugal. – *Acta Biol. Szeged.* 21, 43–62.
- KEDVES, M. and PÁRDUTZ, Á. (1983): Electron microscope investigations of the Early *Normapolles* pollen genus *Atlantopollis*. – *Palynology* 7, 153–169.
- KEDVES, M. et PITTAU, P. (1979): Contributions à la connaissance des pollens des *Normapolles* du "groupe papilloïde" du Crétacé supérieur du Portugal. – *Pollen et Spores* 21, 169–209.
- KRUTZSCH, W. (1959): Einige neue Formgattungen und – Arten von Sporen und Pollen aus der Mitteleuropäischer Oberkreide und dem Tertiär. – *Palaeontographica B*, 105, 125–157.
- KRUTZSCH, W. (1967): In GÖCZÁN, F., GROOT, J. J., KRUTZSCH, W. und PACLTÓVÁ, B.
- KRUTZSCH, W. (1971): Atlas der mittel- und jungtertiären dispersen Sporen- und Pollen- sowie der Mikroplanktonformen des nördlichen Mitteleuropas. Lief. VI Coniferenpollen (*Saccites* und "*Inaperurates*"). – VEB Gustav Fischer Verlag, Jena.
- MARTINS, U. P. (1988): Die Kompressions- und Regressionsphase der obersten Kreide im südlichen Basko-Kantabrischen Becken, Nordspanien (Fazies, Paläogeographie, Paläoenvironment). – Diss. Geowiss. Fak. Eberhard-Karls-Universität Tübingen.
- MARTINS, U. P. (1989): Sedimentary Evolution of the Shallow-Marine Late Cretaceous in the Southern Basco-Cantabrian Basin (Northern Spain). – In: WIEDMANN, J., ed. Cretaceous of the Western Tethys. Proceedings 3rd International Cretaceous Symposium, Tübingen 1987, E. Schweizerbart'sche Verlagsbuchhandlung, Stuttgart, 121–144.
- MÉDUS, J. (1970a): Première contribution à la palynologie stratigraphique du Crétacé supérieur du Sud-Est de la France et du Nord-Est de l'Espagne. – *Paleobiol. Cont.* 1, 3–15.
- MÉDUS, J. (1970b): Contribution à la connaissance des associations polliniques du Crétacé terminal dans le S.E. de la France et le N.E. de l'Espagne. – *Rev. Micropal.* 13, 45–50.
- MÉDUS, J. (1972): Palynological zonation of the Upper Cretaceous in Southern France and Northeastern Spain. – *Rev. Palaeobot. Palynol.* 14, 287–295.
- MÉDUS, J. (1975): Quelques palynoflores du Tertiaire inférieur du Sud de la France et du Nord de l'Espagne. – *Rev. Esp. Micropaleont.* 7, 113–126.
- MÉDUS, J. (1978): Palynological correlation in the Paleocene-Eocene of Southern France and North-Eastern Spain. – *Palinologia* 1, 341–353.
- MÉDUS, J. (1987): Analyse quantitative des palynoflores du Campanien de Cedano, Espagne. – *Rev. Palaeobot. Palynol.* 51, 309–326.
- MÉDUS, J. et ALVAREZ-RAMIS, C. (1989): Du pollens *Normapolles* d'un niveau du Crétacé supérieur de la Region de Guadalix, Prov. de Madrid, Espagne. – *Rev. Esp. de Micropaleont.* 21, 139–144.
- NORRIS, G. (1967): Spores and pollen from the Lower Colorado Group (Albian? Cenomanian) of Central Alberta. – *Palaeontographica B*, 120, 72–115.
- PFLUG, H. D. (1953): Zur Entstehung und Entwicklung des angiospermiden Pollens in der Erdgeschichte. – *Palaeontographica B*, 95, 60–171.
- POCOCK, S. A. J. (1962): Microfloral analysis and age determination of strata at the Jurassic-Cretaceous boundary in the western Canada plains. – *Palaeontographica B*, 111, 1–95.
- PORTA DE, J., KEDVES, M., SOLÉ DE PORTA, N. y CIVIS, J. (1985): Palinología del Maastrichtiense del Barranco de la Posa (Lérida, España). Problemática regional. – *Rev. Inv. Geol.* 40, 5–28.

- SOLÉ DE PORTA, N. (1978): Palinología de dos secciones del Cenomaniense de Los Alrededores de Oviedo (España). – *Palinología*, num. extraord. 1, 435–441.
- TREVISAN, L. (1967): Pollini fossili del Miocene superiore nei tripoli del Gabbro (Toscana). – *Palaeontographia Italica* 62, 1–78.
- TSCHUDY, R. H. (1973): *Complexiopollis* Pollen Lineage in Mississippi Embayment Rocks. – Contribution to Paleontology, Geol. Survey Prof. Paper 743–C, C1–15.
- VAN HOEKEN-KLINKENBERG, P. M. J. (1964): A palynological investigation of some Upper Cretaceous sediments in Nigeria. – *Pollen et Spores* 6, 209–231.

Plate 3.1.

- 1,2. *Stereisporites aegypticus* KEDVES 1986, *Sphagnaceae*, slide: 268/14-1, cross-table number: 8.8/135.3.
3. *Cicatricosisporites baconicus* H. DEÁK 1963, *Schizaeaceae*, slide: 268/14-4, cross-table number: 10.5/135.8.
4. *Appendicisporites potomacensis* BRENNER 1963, *Schizaeaceae*, slide: 268/14-3, cross-table number: 10.3/138.8.
5. Cf. *Clavatisporites* fsp., slide: 268/14-3, cross-table number: 26.2/130.1.
- 6,7. *Liburnisporites* fsp., slide: 268/14-1, cross-table number: 8.4/131.6.
- 8,9. *Camarozonosporites insignis* NORRIS 1967, *Lycopodiaceae*, slide: 268/14-3, cross-table number: 10.9/143.6.
- 10,11. *Ischiosporites* fsp. cf. *tuberosus* DÖRING 1965, slide: 268/14-2, cross-table number: 6.5/127.6.
12. *Ischiosporites* cf. *pseudoreticulatus* (COUPER 1958) DÖRING 1965, slide: 268/14-1, cross-table number: 13.4/127.4.
13. *Uvaesporites* cf. *neerlandicus* HERNGREEN, FELDER, KEDVES and MEESSEN 1986, slide: 268/14-4, cross-table number: 11.5/134.8.
- 14,15. *Podocarpidites multesimus* (BOLKHOVITINA 1956) POCKOCK 1962, slide: 268/14-3, cross-table number: 8.6/130.6.
- 16,17. *Classoidites glandis* AMEROM 1965, slide: 268/14-3, cross-table number: 8.3/129.4.
- 18,19. *Atlantopollis microreticulatus* KRUTZSCH 1967, slide: 268/14-3, cross-table number: 8.3/129.4.
- 20,21. *Atlantopollis vilaflorensis* KEDVES and DINIZ 1979, slide: 268/14-2, cross-table number: 24.2/131.8.
- 22,23. *Atlantopollis grooti* KEDVES and DINIZ 1979, slide: 268/14-1, cross-table number: 15.6/141.7.
24. *Atlantopollis choffatii* DINIZ, KEDVES and SIMONCSICS 1974, slide: 268/14-1, cross-table number: 7.4/137.2.
- 25,26. *Complexiopollis praeatumesens* KRUTZSCH 1959, slide: 268/14-3, cross-table number: 9.8/139.1.
- 27,28. *Complexiopollis praeatumesens* KRUTZSCH 1959, slide: 268/14-4, cross-table number: 17.3/127.4.
- 29,30. *Complexiopollis vulgaris* (GROOT and GROOT 1962) GROOT and KRUTZSCH 1967, slide: 268/14-1, cross-table number: 20.4/143.8.
- 31,32. *Complexiopollis turonis* (KRUTZSCH 1959) KRUTZSCH 1967, slide: 268/14-2, cross-table number: 20.2/121.8.
- 33,34. *Complexiopollis latis* (KRUTZSCH 1959) KRUTZSCH 1967, slide: 268/14-1, cross-table number: 21.3/128.9.

Plate 3.2.

- 1,2. *Pinuspollenites spherisaccus* BRENNER 1963, slide: 194/10-2, cross-table number: 6.7/127.1.
- 3,4. *Podocarpidites verrucatus* (TREVISAN 1967) KRUTZSCH 1971, *Podocarpaceae*, slide: 194/10-1, cross-table number: 10.3/135.3.
- 5,6. *Retitricolpites minutus* (BRENNER 1963) DETTMANN 1973, slide: 194/10-1, cross-table number: 19.9/130.3.
- 7,8. *Ilexpollenites erdmani* KEDVES 1978, *Aquifoliaceae*, slide: 194/10-1, cross-table number: 6.8/138.7.
- 9,10. *Verruculopollis* fsp., slide: 194/10-1, cross-table number: 8.3/130.7.
- 11,12. *Endopollis* fsp., slide: 194/10-2, cross-table number: 20.2/125.9.

- 13,14. *Kruttschippollis spatiosus* GÓCZÁN 1967, slide: 194/10-1, cross-table number: 11.2/127.4.
 15,16. *Semioculopollis medius* KEDVES and DINIZ 1983, slide: 194/10-1, cross-table number: 10.3/130.3.
 17,18. *Papillopollis* fsp., slide: 194/10-2, cross-table number: 18.2/127.2.

Plate 3.3.

- 1,2. *Papillopollis pflugii* KEDVES and PITTAU 1979, slide: 194/10-2, cross-table number: 12.9/129.3.
 3,4. *Papillopollis vancampoae* KEDVES and PITTAU 1979, slide: 194/10-2, cross-table number: 22.2/140.8.
 5,6. *Papillopollis vancampoae* KEDVES and PITTAU 1979, slide: 194/10-1, cross-table number: 18.4/135.2.
 7,8. *Papillopollis rugulatus* KEDVES and PITTAU 1979, slide: 194/10-1, cross-table number: 6.8/136.9.
 9,10. *Papillopollis regulus* PFLUG 1953, slide: 194/10-1, cross-table number: 17.7/130.9.
 11,12. *Rocheipollenites triangulus* KEDVES and DINIZ 1980, slide: 194/10-2, cross-table number: 9.3/131.6.
 13,14. *Interporopollenites plicatus* KEDVES and HEGEDÜS 1975, slide: 194/10-2, cross-table number: 21.2/128.9.
 15,16. *Triangulipollis* cf. *magnus* KEDVES and DINIZ 1980–1981, slide: 194/10-2, cross-table number: 8.3/129.9.
 17,18. *Interporopollenites plicatus* KEDVES and HEGEDÜS 1975, slide: 194/10-1, cross-table number: 9.2/129.3.
 19,20. *Interporopollenites stanley* KEDVES and HEGEDÜS 1975, slide: 194/10-1, cross-table number: 20.8/136.5.
 21,22. *Interporopollenites concavus* KEDVES and HEGEDÜS 1975, slide: 194/10-2, cross-table number: 24.3/123.4.
 23,24. *Interporopollenites dinizae* KEDVES and HEGEDÜS 1975, slide: 194/10-1, cross-table number: 16.3/140.6.
 25,26. *Interporopollenites ornatus* KEDVES and HEGEDÜS 1975, slide: 194/10-2, cross-table number: 22.3/134.2.
 27,28. *Alnipollenites* fsp., slide: 194/10-1, cross-table number: 16.5/142.2.
 29,30. *Triporopollenites nointelensis* KEDVES 1970, *Corylaceae*, slide: 194/10-1, cross-table number: 12.2/136.1.
 31,32. *Triatriporopollenites lubomirovae* (GLADKOVA 1965) KEDVES 1974, *Myricaceae*, slide: 194/10-2, cross-table number: 9.9/136.9.
 * *Complexiopollis* cf. *paulus* TSCHUDY 1973.
 33,34. *Inaperturotetradites lacunosus* VAN HOEKEN-KLINKENBERG 1964, slide: 194/10-2, cross-table number: 17.6/141.6.




Plate 3.1.




Plate 3.2.




Plate 3.3.

4. CONSIDERACIONES EN TORNO A LAS ASOCIACIONES ESPOROPOLINICAS DEL CRETACICO SUPERIOR DEL CERRO DE LA MESA (GUADALIX DE LA SIERRA, MADRID)

C. ALVAREZ RAMIS¹, M. KEDVES² y T. FERNÁNDEZ MARRÓN¹

1. Departamento de Paleontología Facultad de Ciencias Geológicas (U. C. M.), 28040 Madrid / U. E. I. de Paleontología Inst. Geología Económica C. S. I. C. Madrid, España, 2. Laboratório de Biología celular y Micropaleontología evolutiva, Departamento de Botánica, Universidad J. A., H-6701, Apartado Postal 657, Szeged, Hungria

Resumen

Se discuten los resultados obtenidos del estudio paleopalinológico efectuado en 7 niveles, tomados a lo largo de una serie litoestratigráfica establecida en sedimentos cretácicos de la provincia de Madrid. Se determinan los porcentajes de los principales componentes de la paleoasociación y se comparan con los de otros yacimientos cretácicos, más o menos sincrónicos

Palabras clave: Palinología, fósil, Cretácico superior, provincia de Madrid, España.

Introducción


Este estudio forma parte de una serie de proyectos que nos han sido concedidos ininterrumpidamente por el CSIC y la DG CYT desde 1982 hasta 1995 entre los que se incluye un proyecto concedido por la comisión mixta Hispano-Hungara para 1992 y 1993.

Las muestras estudiadas se han tomado en diferentes puntos de los tramos 4 y 6 de la serie estratigráfica local figs. 4.2., 4.3., 4.4. y 4.5. siendo todas ellas ricas en palinomorfos (cutículas, leños, algas, esporas y sobre todo pólenes) además de algunas cámaras internas de *Foramíniferos*.

Localización y acceso al yacimiento

El yacimiento del Cerro de la Mesa se encuentra en la Comunidad Autónoma de Madrid. Geográficamente se situa en el borde meridional de la Sierra de Guadarrama que forma parte del Sistema Central.

Los materiales estudiados proceden de una explotación, a cielo abierto, de la que se extraen bloques de caliza compacta empleados en la construcción. La cantera se situa en un cerro de cima plana, mesa, ubicado en una zona próxima al pueblo de Guadalix de la Sierra.



- A** ANGIOSPERMAS PRIMITIVAS
- B** GIMNOSPERMAS
- C** ESPORAS
- D** ANGIOSPERMAS TIPO ACTUAL
- E** OTROS

Figura 4.1.

El emplazamiento preciso de la columna establecida en el Cerro de la Mesa corresponde a las coordenadas (Greenwich) 40° 47' 37" N y 3° 42' 05" W.

El acceso al yacimiento, desde la ciudad de Madrid, se efectúa a través de la Carretera Nacional I hasta el kilómetro 50 en que ésta se cruza, casi perpendicularmente, con las carreteras que conducen a Torrelaguna (por el Este) y a Guadalix de la Sierra (por el Oeste). Siguiendo por esta última se llega al pueblo de Guadalix de la Sierra, donde se toma, en dirección Norte, otra carretera que conduce a Navalafuente; de ella parte el camino que da servicio a varias explotaciones ganaderas y que conduce a la cantera del Cerro de la Mesa.

Aspectos geológicos

La zona en que se emplaza el yacimiento forma parte del Sistema Central y se sitúa en el borde meridional de la Sierra de Guadarrama. Los materiales cretácicos, junto con los paléogenos rellenan una cubeta que se dispone en dirección SW-NE formando la cobertera de la Sierra de Guadarrama (ALVAREZ-RAMIS, FERNÁNDEZ MARRÓN y GOMEZ PORTER, 1984, CORCHON RODRIGUEZ, 1976 y FUSTER y FEBREL, 1959). En la zona en que se sitúa el yacimiento el Cretácico forma un sinclinal que cierra periclinalmente en dirección al pueblo de Guadalix de la Sierra y cuyo eje se encuentra más próximo al flanco norte. El Cerro de la Mesa se sitúa en el flanco sur del Sinclinal; los materiales cretácicos que integran sus capas presentan un buzamiento de 45° N. Los materiales que constituyen los niveles cretácicos están formados fundamentalmente por calizas (que en algunas capas pueden presentar elevados porcentajes de margas o arenas) y otros niveles de arcillas de escasa potencia. La disposición de los materiales se caracteriza por presentar una alternancia cíclica. La sucesión litológica, de donde proceden las muestras, es la característica de todas las canteras situadas en el Borde Sur de la Sierra de Guadarrama. La serie local establecida, está formada por calizas más o menos compactas (algunas con porcentajes elevados de arena) y margas. De forma discontinua aparecen intercalados pequeños niveles carbonosos (figuras 4.2., 4.3., 4.4. y 4.5.). La columna forma parte de una de las unidades litoestratigráficas establecidas por CORCHON RODRIGUEZ en 1976.

Caracterización palinológica del Cretácico superior del Borde Sur de la Sierra de Guadarrama

La inclusión en este artículo de este apartado se justifica en base a que para hacer una caracterización fiable de paleobioasociaciones, es necesario conocer previamente las características de otros yacimientos de la misma edad, o de niveles próximos, enclavados en el área en que se localiza el yacimiento. Esta circunstancia se cumple en este caso, ya que en el Cretácico superior del borde meridional de la Sierra de Guadarrama se han estudiado otros yacimientos, la mayoría de ellos en profundidad. Al haberse iniciado, en el último año, el reconocimiento de tres nuevos yacimientos, nos hace variar ligeramente los porcentajes del conjunto de los distintos grupos sistemáticos dados a conocer en el 8° Simposio de Palinología (ALVAREZ RAMIS, y FERNÁNDEZ MARRÓN, 1990). No hemos estimado necesario dar a conocer aisla-




Figura 4.2.

damente los porcentajes de cada nivel ya que aunque las preparaciones pueden presentar entre si marcadas diferencias, sus promedios son muy semejantes tanto en porcentajes como en tipos botánicos. Los grupos establecidos fueron: Esporas, pólenes de *Gymnospermas*, pólenes de *Angiospermas* primitivas, pólenes de *Angiospermas* de tipo actual y otros. Este apartado carece de rigor paleobiológico, al incluir representantes del reino *Proctista* (*Dinoflagelados*, *Botryococcales*), epidermis vegetales, vasos conductores, etc. junto a epidermis de animales, cámaras internas de *Foraminíferos* etc. Seguidamente indicamos, en porcentajes, los componentes comunes que hemos establecido en todos los yacimientos estudiados (figura 4.1.).


Aspectos paleopalinológicos del Cerro de la Mesa

En 1990 ALVAREZ RAMIS y FERNÁNDEZ MARRÓN dimos a conocer el estudio palinológico preliminar de este yacimiento, indicando el promedio de los distintos porcentajes obtenidos en los diversos niveles de la serie establecida. La metodología empleada en la obtención de los palinomorfos fué la citada en publicaciones anteriores por los autores para los grana incluidos en niveles de carbones y margas. En la continuación de ese estudio, que damos a conocer aquí, se han mantenido en principio los mismos grupos básicos establecidos, si bien a la vista de los resultados que se iban obteniendo se profundizó en el conocimiento de las *Angiospermas* por su importancia e interés en aspectos tales como la filogenia, evolución, áreas de dispersión y valor estratigráfico de estas plantas durante el Cretácico superior. No pormenorizamos los distintos géneros sistemáticos obtenidos en el estudio de las paleoasociaciones del yacimiento aunque se han reconocido nuevas formas de *Dinoquistes* y sobre todo de *Normapólenes* que constituyen el objetivo básico de esta publicación. Los valores que arrojan los nuevos porcentajes obtenidos se desvían poco de los que habíamos obtenido previamente, siendo los más significativos el aumento de las *Gymnospermas* (5%) y la disminución de las *Criptógamas* (4%). Detallando los porcentajes determinados en los siete niveles en que se ha realizado el muestro los resultados obtenidos ponen de manifiesto que las *Angiospermas* son dominantes en todos ellos, seguidos muy se cerca por los pólenes de *Gymnospermas*, siendo casi idénticos en el tramo 6.


Estos porcentajes se expresan gráficamente a la derecha de la serie local establecida (figura 4.2.).

En otra figura puede apreciarse la distribución en la serie de los porcentajes hallados para los tipos polínicos más representativos de *Coníferas* (figura 4.3.). Los porcentajes y distribución obtenidos para el Cerro de la Mesa no son concordantes con los obtenidos en yacimientos sincrónicos estudiados en la zona. (ALVAREZ RAMIS 1985, ALVAREZ RAMIS y GOMEZ PORTER, 1985 y ALVAREZ RAMIS y FERNÁNDEZ MARRÓN 1990).

La siguiente figura (figura 4.4.) se refiere a las *Angiospermas*, indicando los porcentajes obtenidos para los distintos niveles en que se han estudiado los palinomorfos. En el centro de la figura se representan los porcentajes de *Angiospermas* cuyo eje polar supera al eje ecuatorial. Este tipo lo presentan con mayor frecuencia las *Angiospermas* actuales. A la derecha de la figura se representan los pólenes *Breviejes*



10%



Classopollis

Tipo Taxodiaceae/
Cupressaceae

Araucariacites

Disacados

Figura 4.3.




Figura 4.4.

que son los típicos de las *Angiospermas* primitivas. Estas formas caracterizan el Cretácico superior, época en que destacan por su cantidad y diversidad los *Normapólenes*. Como puede observarse en esta figura todos los niveles presentan una gran desproporción entre los porcentajes de los pólenes *Brevaxones* y *Longaxones* en favor de los primeros, a excepción del nivel intermedio del tramo 4 en que ambos son muy semejantes. Dado el interés que representan las formas *Breviejes* en las dataciones estratigráficas y correlaciones del Cretácico superior se representan en la figura 4.5., porcentajes de los tipos polínicos cuya presencia en los distintos niveles muestreados ha resultado significativa (*Interporopollenites*, *Papillopollis*, *Complexiopollis* y *Atlantipollis*).

Discusión de resultados

Las especies y formas esporopolínicas halladas en el yacimiento del Cerro de la Mesa son, en líneas generales, las mismas que hemos determinado en los otros




Figura 4.5.



yacimientos estudiados en el Cretácico superior del Borde Sur de la Sierra de Guadarrama.

No obstante la presencia o abundancia de formas con valor cronoestratigráfico, especialmente palinomorfos pertenecientes a *Angiospermas* primitivas y/o a quistes de *Dinoflagelados* nos permite especular con más precisión acerca de la posición estratigráfica de los niveles estudiados. La dominancia, entre los *Normapólenes*, de *Interporopollenites*, así como la variedad de tipos de *Papillopollis* y de *Dinoquistes* (estos serán estudiados posteriormente por un especialista en la materia) entre los que creemos reconocer diversos tipos de *Deflandrea*, *Weitzellites* y sobre todo la abundancia de *Mycrodinium* afín a *M. veligerum* o a *M. kustanaikum* que son formas que caracterizan el Campaniense y Maastrichtiense, parecen datar los niveles estudiados.

Si se comparan las asociaciones encontradas en el Cerro de la Mesa con las de otros yacimientos del borde meridional de la Sierra de Guadarrama no se observan diferencias notables, como ya se ha indicado anteriormente.

En relación a la palinoflora descrita en Torrelaguna (ALVAREZ RAMIS y DOUBINGER, 1980) existen numerosas formas comunes, sobre todo las más recientes. El yacimiento fué atribuido al intervalo sedimentario Cenomaniense superior-Santonienense. Si se compara con otros yacimientos más o menos sincrónicos de la Península Ibérica, aparecen tipos polínicos semejantes a los descritos en Vila Flor (Cenomaniense superior-Turonienense) y Carrajão (Turonienense-Coniacienense) por DINIZ, KEDVES y SIMONCSICS en 1974 y en Aveiro (Santonienense-Campanienense) por KEDVES y DINIZ en 1967. Hay que destacar en Cerro de la Mesa la frecuencia relativa de *Rocheipollenites triangulus* y la presencia de *Aveiropollenites triangulus* descritos por primera vez en Portugal y escasos en la localidad tipo.

La coincidencia entre los palinomorfos hallados en nuestro yacimiento y la de los que integran las diferentes paleoasociaciones cretácicas de la Península Ibérica, junto con la abundancia de *Interporopollenites* y *Papillopollis*, presencia de formas triporadas de tipo actual y de *Dinoflagelados* que caracterizan las series finicretácicas (Campanienense-Maastrichtiense) parecen reforzar la datación propuesta para los niveles muestreados en el Cerro de la Mesa como al menos Campanienense.

La iconografía de los *Normapólenes* de mayor interés se ha figurado, por los autores en Plant Cell Biology and Development 4, 22–25, y un aspecto general de la paleoasociación se da a conocer en la lámina del trabajo presentado al 9º Simposio de Palinología; Las Palmas de Gran Canaria – España 1992.

Bibliografía

- ALVAREZ RAMIS, C. (1980): Sur la macroflore du Crétacé continental de l'Espagne. – Mem. Soc. Géol. France, N. S. 139, 5–9.
- ALVAREZ RAMIS, C. (1981): Estudio de *Frenelopsis oligostomata* (ROMARIZ) ALVIN, procedente del Cretácico superior de Torrelaguna (Madrid). – Estudios Geol. 37, 83–87.
- ALVAREZ RAMIS, C. (1985): Sur la paléocologie des alentours du Barrage "El Vellón" (Madrid, Espagne) pendant le Crétacé supérieur. – Sciences (M. E. N. Paris) 5, 115–125.
- ALVAREZ RAMIS, C. y DOUBINGER, J. (1980): Los microfósiles vegetales del Cretácico superior de Torrelaguna (Madrid). – Bol. Real Soc. Española de Historia Natural (Geol.) 78, 205–215.

- ALVAREZ RAMIS, C. y DOUBINGER, J. (1989): Contribution à l'étude de la palynoflora Crétacée de Guadalix de la Sierra (Madrid, Espagne). – II European Paleobotanical Conference Abstracts, 15.
- ALVAREZ RAMIS, C. y FERNÁNDEZ MARRÓN, T. (1990): Palinomorfos atribuidos a Gimnospermas hallados en niveles supracretácicos del embalse de "El Vellón" (Provincia de Madrid). – Polen, Esporas y sus aplicaciones. VII Simposio de Palinología A. P. L. E. Granada, 437–443.
- ALVAREZ RAMIS, C. y FERNÁNDEZ MARRÓN, T. (en prensa): Caracterización del Cretácico superior del Cerro de la Mesa (Guadalix de la Sierra, Madrid). – 8º Simposio de Palinología A. P. L. E. Tenerife, 1990.
- ALVAREZ RAMIS, C., FERNÁNDEZ MARRÓN, T. y GOMEZ PORTER, P. (1984): Etude des niveaux à plantes du Crétacé supérieur d'une carrière située au bord du Barrage "El Vellón" (Madrid, Espagne). – Sciences (M. E. N.) II, 83–97.
- ALVAREZ RAMIS, C., FERNÁNDEZ MARRÓN, T. y GOMEZ PORTER, P. (1989): Avance del estudio macropaleobotánico de la flora supracretácica del Cerro de la Mesa (Sierra de Guadarrama, Madrid). – II European Paleobotanical Conference Abstracts, 15.
- ALVAREZ RAMIS, C. y GOMEZ PORTER, P. (1985): Nouvelles données concernant le Crétacé supérieur de la provincia de Madrid (Espagne). Carrière de la "Calerizas." – Sciences (M. E. N. Paris) V, 137–146.
- ALVAREZ RAMIS, C. y GOMEZ PORTER, P. (1990): Estudio palinológico de Pteridófitos del Cretácico superior de la provincia de Madrid. 1ª parte: Embalse de El Vellón. – Acta Salamanticensis 68, 47–53.
- ARIAS ORDÁS, C. (1969): Estudio estratigráfico de los alrededores de Guadalix de la Sierra (Madrid). – Cuadernos de Geología Ibérica I, 309–333.
- CORCHON RODRIGUEZ, F. (1976): Estudio hidrogeológico del Cretácico de los alrededores de Torrelaguna (Madrid y Guadalajara). – Informaciones y estudios, Servicio Geológico. M. O. P.
- DINIZ, F., KEDVES, M. et SIMONCSICS, P. (1974): Les sporomorphes principaux de sédiments crétacés de Vila Flor et de Carrajão. – Com. Serv. Geol. de Portugal 58, 161–178.
- FUSTER, J. M. y FEBREL, T. (1959): Hoja nº 509. Torrelaguna. Mapa Geológico de España IGME 1:50.000.
- KEDVES, M. (1981): Definitions of evolutionary trends within and classifications of Early Brevaxonate pollen. – Rev. Palaeobot. Palynol. 35, 149–154.
- KEDVES, M., ALVAREZ RAMIS, C. y FERNÁNDEZ MARRÓN, M. T. (1993): Sobre polenes Breviaxones procedentes del Crétacico superior del borde sur de la Sierra de Guadarrama (Provincia de Madrid, España). – Plant Cell Biology and Development 4, 22–25.
- KEDVES, M. et DINIZ, F. (1967): Quelques types de sporomorphes de sédiments crétacés d'Aveiro, Portugal. – Com. Serv. Geol. de Portugal 52, 17–25.
- KEDVES, M. et DINIZ, F. (1979): Les pollens d'*Angiospermes* du Crétacé de Vila Flor, Portugal, Genres de forme *Atlantopollis* et *Limaipollenites*. – Bol. Soc. Geol. de Portugal 21, 203–216.
- KEDVES, M. et DINIZ, F. (1981): Contribution à la connaissance des pollens d'*Angiospermes* du Crétacé supérieur du Portugal. – Bol. Soc. Geol. de Portugal 22, 19–36.
- KEDVES, M. et PITTAU, P. (1979): Contribution à la connaissance de pollens des *Normapollis* du Groupe *Papilloide* du Crétacé supérieur du Portugal. – Pollen et Spores 21, 169–209.
- MÉDUS, J. et ALVAREZ RAMIS, C. (1989): Des pollens *Normapollis* d'un niveau du Crétacé supérieur de la région de Guadalix, Prov. de Madrid, Espagne. – Revista española de Micropaleontología 20, 139–144.
- PORTA, J., KEDVES, M., SOLÉ DE PORTA, N. y CIVIS, J. (1985): Palinología del Maastrichtense del Barranco de la Posa (Lérida, España). Problemática regional. – Rev. Inv. Geol. 40, 5–28.
- SOLÉ DE PORTA, N. (1978): Palinología de dos secciones del Cenomaniense de los alrededores de Oviedo (España). – Palinología. N° extraordinario I, 435–441.

5. TEM STUDY OF THE "INFECTED" POLLEN GRAINS OF *THALICTRUM FLAVUM* L. WITH *GLIOCLADIUM ROSEUM* (LINK) BAINIER

Short communication

M. KEDVES¹ and I. ROJIK²

1. Cell Biological and Evolutionary Micropaleontological Laboratory of the Department of Botany of the J. A. University, H-6701, P. O. Box 657, Szeged, Hungary, 2. Faculty of Science, Laboratory of Electron-microscopy, P. O. Box 553, Szeged, Hungary

During our experimental investigations of the secondary alterations of the spores and pollen grains (e. g.: KEDVES and KINCSEK, 1989, KEDVES and PÁRDUTZ, 1992a), several taxa were examined. The pollen grains of the interesting, wind-pollinated *Ranalean* genus – *Thalictrum* – were also the subjects of our investigations. On 17. 6. 1988, 20 mg air dried pollen material was measured six times in small glasses for high temperature effect experiments. Not long after the preparation of the experiment, symptoms of infections were observed. The infected pollen grains were placed into a thermostat at 30 °C for 24^{hrs}. The determination of the microscopical fungi was made in the Department of Microbiology of the J. A. University, by Dr. L. MANCZINGER, as *Gliocladium roseum* (LINK) BAINIER. The infected material was fixed (OsO₄ aq. dil. 1%), embedded in Araldite, ultrathin sectioned, and investigated with the TEM method. The electron microscopical pictures were taken with a Tesla BS-500 transmission electron microscope; resolution: 6 Å. Our preliminary results can be summarized as follows.


1. Fungal cells, mostly hyphae were observed intramatically (inside the pollen grain) and extramatically (outside the pollen grain); cf. HESSE, KUSEL-FETZMANN and CARNIEL, (1989) (Plate 5.1., plate 5.2., figs. 1,2).
2. Intramatical fungal cells can grow through the aperture of the pollen grain, and this kind of hypha is similar to the pollen tube; Plate 5.1., marked with two arrows.

Plate 5.1. ►

Thalictrum flavum L. Recent pollen grain with *Gliocladium roseum* (LINK) BAINIER.

The fungal cells are shown in the protoplasm of the pollen grains and outside the exine. It is worth mentioning that several fungal cells reach the plasma membrane of the pollen grain. Marked with an arrow. One of them is growing out through the pollen aperture pushing out the thin exine layer of the germinal area. Indicated with two arrows. The thin electron dense coat layer on the surface is well shown. Extramatical fungal cells may be in close contact with the surface of the pollen grain.

Negative no: 8946, 10.000 x.



3. Intramatrix fungi can be nourished from the protoplasm, completely empty exines were also observed. Cf.: FAEGRI (1971), BROOKS and SHAW (1973).

4. Extramatrix fungi can also digest the remains of the tapetal tissues.

5. The interactions between the surface of the pollen grain and the fungal cells are important. Fungal cells can grow directly towards the pollen grain. This can be the consequence of several factors, as follows.

5.1. The peculiar biopolymer system of the pollen surface (cf. KEDVES, PÁRDUTZ, and VÉR, 1991, KEDVES and PÁRDUTZ, 1992b).

5.2. The electrostatic charge of the pollen surface (ROWLEY, 1971), namely the anionic character, which results in the accumulations of the cations, particularly the thorium on the surface.

5.3. The peculiar characteristics of the pollen coat (tryphine). Several concepts were published concerning this question; EHRLICH (1958), FREYTAG, (1967), SKVARLA and ROWLEY (1970), DICKINSON and LEWIS (1973), HESLOP-HARRISON (1975), MASCARENHAS (1975), GILLISSEN and BRANTJES (1978), HESSE (1978a, b, c, 1979, 1980, 1984, 1991), AUDRAN and BATCHO (1981), ZAVADA (1983), BLACKMORE and BARNES (1984), LOEWUS et al. (1985), SOUTHWORTH (1985), FERNÁNDEZ and RODRIGUEZ-GARCIA (1988), HARLEY (1988, 1991), DICKINSON and SHELDON (1990), VAN DER HAM (1990), ROMAN, BLANCA and ROMERO (1992). The chemical interaction between the surfaces of the fungal cell and the pollen grain is also a factor in this phenomenon.

5.4. The molecular sieving character of the exine (ROWLEY, 1973) is also important and so is the basic concept of ROWLEY (1987–88) which says that the “pollen wall” is an integrated part of the plasma membrane.



5.5. The plasma membrane and the tapetal membrane glycolyx are also important in this interaction. The fungal cell, which is in connection with the plasma membrane (glycolyx), can “imitate” the pollen tube.

6. According to the newest concept of fungal-plant interactions, the following can be cited from the book of ISAAC (1992), p. 316: “There is presently a great deal of discussion concerning the extent to which endophytes are in fact latent pathogens of examples of fungi co-evolving with plants from parasitism to mutualism (CLAY, 1988). However in many cases mutualistic relationships between host plants and fungi endophytes are suspected and in many instances such associations have been identified.” P. 320. “Many endophytes live almost entirely within the host plant tissues, often without causing any visible signs of infection.”

Plate 5.2. ►

Thalictrum flavum L. Recent pollen grain with *Gliocladium roseum* (LINK) BAINIER.

1. Detail of the pollen exine ultrastructure with a fungal cell on the surface. The contact between the pollen and fungal cell surfaces is very close in some parts. The disintegration (digestion by fungal enzymes) of the pollen coat is well shown. negative no: 8614, 40.000x.
2. Detail of the exine with remains of the tapetal tissues. One fungal cell is growing directly towards the pollen exine. On the tectum, the thin electron dense layer, the pollen coat is well illustrated. Negative no: 8589, 20.000x.



In resumé, the observed fungal-pollen interaction is a complicated, multifactorial biochemical and physiological relationship.

Finally it is necessary to emphasize that to get more information about the relationship of the connected surfaces of the pollen and the fungal cell, we need further TEM data with an instrument of better resolution power. These investigations are in progress, and the new results will be published in the near future.

The authors are deeply indebted to Prof. DR. L. FERENCZY, member of the Hungarian Academy of Sciences, and to DR. L. MANCZINGER, first assistant (Department of Microbiology of the J. A. University, Szeged, Hungary) for their valuable comments and advice.

This work was supported by the Grant OTKA 1/3-104.

References

- AUDRAN, J.-C. and BATCHO, M. (1981): Cytochemical and infrastructural aspects of pollen and tapetum ontogeny in *Silene dioica* (Caryophyllaceae). – *Grana* 20, 65–80.
- BLACKMORE, St. and BARNES, S. H. (1984): Transmission electron microscope preparations from unacetolysed herbarium pollen: A comparison of methods. – *Pollen et Spores* 26, 137–146.
- BROOKS, J. and SHAW, G. (1973): The role of sporopollenin in Palynology. In: *Problems of Palynology*, Nauka, Moscow, 80–91.
- DICKINSON, H. G. and LEWIS, D. (1973): The formation of the tryphine coating the pollen grain of *Raphanus* and its properties relating to the self-incompatibility systems. – *Proc. R. Soc. (London)* 183, 149–165.
- DICKINSON, H. G. and SHELDON, J. M. (1990): The cell biological basis of exine formations in *Lilium* sp. – *Polen esporas y sus aplicaciones*, VII Simposio de Palinologia, 17–29.
- EHRlich, H. G. (1958): Electron microscope studies of *Saintpaulia ionantha* WENDL. pollen walls. – *Experimental Cell Research* 15, 463–474.
- FAEGRI, K. (1971): The preservation of sporopollenin membranes under natural conditions. In: *Sporopollenin* eds.: J. BROOKS, P. R. GRANT, M. MUIR, P. VAN GUZEL and G. SHAW. – Academic Press, London, New York, 256–272.
- FERNÁNDEZ, M. C. and RODRIGUEZ-GARCIA, M. I. (1988): Pollen wall development in *Olea europaea* L. – *New Phytol.* 108, 91–99.
- FREYTAG, K. (1967): Über doppelbrechende Lipide in der Exine und im Pollenöl. – *Grana Palynologica* 7, 3–9.
- GILLISSEN, L. J. W. and BRANTJES, N. B. M. (1978): Function of the pollen coat in different stages of the fertilization process. – *Acta Bot. Neerl.* 27, 205–212.
- HARLEY, M. M. (1988): Observations on the colours of fresh pollen with the light microscope and macroscope. – *Polen, esporas y sus aplicaciones VII Simposio de Palinologia A. P. L. E. Granada 26/30 Septbre 1988 Programa y Resúmenes*, 31.
- HARLEY, M. M. (1991): The pollen morphology of the *Sapotaceae*. – *Kew Bulletin* 46, 379–491.
- HESLOP-HARRISON, J. (1975): The Cronian Lecture, 1974 The physiology of the pollen grain surface. – *Proc. R. Soc. London B*, 190, 275–299.
- HESSE, M. (1978a): Der Feinbau der Pollenklebstoffe: Preparative Probleme bei der Strukturhaltung, Grundfragen zur Nomenklatur und zur Begriffsabgrenzung. – *Linzer Biol. Beitr.* 9, 181–201.
- HESSE, M. (1978b): Entwicklungsgeschichte und Ultrastruktur des Pollenkitts bei *Tilia* (Tiliaceae). – *Pl. Syst. Evol.* 120, 13–30.
- HESSE, M. (1978c): Entwicklungsgeschichte und Ultrastruktur von Pollenkitt und Exine bei nahe verwandten entomophilen und anemophilen Angiospermensippen: *Ranunculaceae*, *Hamamelidaceae*, *Platanaceae* und *Fagaceae*. – *Pl. Syst. Evol.* 130, 13–42.
- HESSE, M. (1979): Entwicklungsgeschichte und Ultrastruktur von Pollenkitt und Exine bei nahe verwandten entomophilen und anemophilen Sippen der *Oleaceae*, *Scrophulariaceae*, *Plantaginaceae* und *Asteraceae*. – *Pl. Syst. Evol.* 132, 107–139.

- HESSE, M. (1980): Ultrastruktur und Entwicklungsgeschichte des Pollenkitts von *Euphorbia cyparissias*, *E. palustris* und *Mercurialis perennis*. – Pl. Syst. Evol. 135, 253–263.
- HESSE, M. (1984): Pollenkitt is lacking in *Gnetatae: Ephedra* and *Welwitschia*: Further proof for its restriction to the *Angiosperms*. – Pl. Syst. Evol. 144, 9–16.
- HESSE, M. (1991): Cytology and Morphogenesis of Pollen and Spores. – Progress in Botany 52, 19–34.
- HESSE, M., KUSEL-FETZMANN, E. and CARNIEL, J. (1989): Life cycle and ultrastructure of *Ducellieria chodati* (*Oomycetes*). – Pl. Syst. Evol. 165, 1–15.
- ISAAC, S. (1992): Fungal-Plant Interactions. – Chapman and Hall, London – New York – Tokyo – Melbourne – Madras.
- KEDVES, M. and KINCSEK, I. (1989): Effect of the high temperature on the morphological characteristic features of the sporomorphs I. – Acta Biol. Szeged. 35, 233–235.
- KEDVES, M. and PÁRDUTZ, Á. (1992a): X-ray effect to the exine ultrastructure of *Alnus glutinosa* (L.) GAERTN. – Taiwania 37, 123–133.
- KEDVES, M. and PÁRDUTZ, Á. (1992b): Transmission electron microscopy of partially dissolved exines of different bisaccate *gymnosperm* pollen grains. In: Plant Cell Biology and Development 3, ed.: M. KEDVES, 38–66, Szeged.
- KEDVES, M., PÁRDUTZ, Á. and VÉR, A. (1991): Biopolymer organization of partially degraded exines of saccate *gymnosperm*. In: Plant cell Biology and Development 1, ed.: M. KEDVES, 32–34, Szeged.
- LOEWUS, F. A., BALDI, B. G., FRANCESCHI, V. R., MEINERT, L. D. and MCCOLLUM, J. J. (1985): Pollen Sporoplasts: Dissolution of Pollen Walls. – Plant Physiol. 78, 652–654.
- MASCARENHAS, J. P. (1975): The biochemistry of *Angiosperm* pollen development. – Bot. Rev. 41, 259–344.
- ROMAN, J. M., BLANCA, G. and ROMERO, A. T. (1992): Consideraciones sobre la presencia de trifina en el polen de *Cruciferas*. – 9 Simposio de Palinologia, A. P. L. E., Resumenes, 86.
- ROWLEY, J. R. (1971): Implications on the nature of sporopollenin based upon pollen development. In: Sporopollenin eds.: J. BROOKS, P. R. GRANT, M. MUIR, P. VAN GUZEL and G. SHAW. – Academic Press. London, New York, 174–219.
- ROWLEY, J. R. (1973): Translocation through the pollen wall. – J. Ultrastructure Research 44, 449–450.
- ROWLEY, J. R. (1987–1988): Substructure within the endexine, an interpretation. – J. of Palynology 23–24, 29–42.
- SKVARLA, J. J. and ROWLEY, J. R. (1970): The pollen wall of *Canna* and its similarity to the germinal apertures of other pollen. – Amer. J. Bot. 57, 519–529.
- SOUTHWORTH, D. (1985): Pollen exine substructure. I. *Lilium longiflorum*. – Amer. J. Bot. 72, 1274–1283.
- VAN DER HAM, R. W. J. M. (1990): *Nephelieae* pollen (*Sapindaceae*): Form, Functions and Evolution. – Proefschrift, Rijksherbarium/ Hortus Botanicus, Leiden.
- ZAVADA, M. S. (1983): Pollen wall development of *Zamia floridiana*. – Pollen et Spores 25, 287–304.

6. BUCKMINSTERFULLERENE-LIKE BIOPOLYMER UNITS FROM THE EXINE OF *THALICTRUM FLAVUM* L.

M. KEDVES¹ and I. ROJIK²

1. Cell Biological and Evolutionary Micropaleontological Laboratory of the Department of Botany of the J. A. University, H-6701, P. O. Box 657, Szeged, Hungary, 2. Faculty of Science, Laboratory of Electron-microscopy, P. O. Box 553, Szeged, Hungary

Abstract

On the TEM micrographs of the partially degraded and fragmented exines of *Thalictrum flavum* L., quasi-equivalent biopolymer structures were observed for the first time from *angiosperm* exines. The diameter of the large globular units is 15 – 35 – 85 Å, and the diameter of the superficial electron dense particles is 5 – 10 – 15 Å. The arrangement of these large globular units may be linear, irregular or they may form network systems of different kinds of polygons. Further peculiarities of the pollen grains of the wind-pollinated *Thalictrum* genus were established experimentally. The quasi-equivalent biopolymer structures were discovered in the partially degraded wall of *Botryococcus braunii* KÜTZ. isolated from oil shale. These structures can be modelled with fullerenes.

Key words: Palynology, recent, *Angiospermae*, quasi-equivalent biopolymer structure.



Introduction

During our investigations on the biopolymer structure and organization levels of the plant cell wall, surprising and unusual biopolymer units were observed on the partially degraded and fragmented wall of *Botryococcus braunii* KÜTZ. extracted from the oil shales of Hungary (KEDVES, ROJIK and VÉR, 1991). The peculiarities of the biopolymer organization of the fossil *Botryococcus braunii* wall were the subjects, among others, of further investigations (KEDVES, ROJIK and VÉR, 1992). The previous-

Plate 6.1. ►

1–4. *Thalictrum flavum* L. Recent.

1. Ultrastructure of the interapertural area of the fresh (non-experimental) pollen grain. Negative no: 8550, 50.000x.
2. Exine ultrastructure of the apertural area of the fresh (non-experimental) pollen grain. Negative no: 8552, 50.000x.
3. Experiment No: 281, globular highly organized biopolymer structures from partially degraded and fragmented exine. Negative no: 9816, 100.000x.
4. Experiment No: 266, detail from the highly organized buckyball-like biopolymer structure. The superficial electron dense molecular systems are well shown. Negative no: 9753, 500.000x.



ly described, large peculiar, and globular units can be modelled with fullerenes having a buckyball-like structure; review of the fullerenes from BECK and BRAUN (1992). Taking into consideration our up-to-date knowledge about chemistry and biopolymer structures, the following scheme can be summarized:

1. Chemical compounds of the plant cell wall.
- 2.1. Basic regular pentagonal biopolymer units of angstrom dimension ($8 - 22 \text{ \AA}$). These units form the quasi-crystalloid metastable skeleton of the plant cell wall in pentagon dodecahedron systems. Such biopolymer systems can be modelled after the theory from PENROSE's tiling patterns (PENROSE, 1979).
- 2.2. In living systems, the metastable quasi-crystalloid biopolymer skeleton is stabilized by another biopolymer system (Cf. KEDVES and TÓTH, 1992, 1994).
- 2.3. The basic regular pentagonal dodecahedron of the PENROSE-I units are the elementary components of the units of a further level of organization in nanometer dimension filaments, lamellae, helical, tubular (ROWLEY et al., 1981), globular (HESSE, 1985), irregular polygons (SOUTHWORTH, 1986), etc.
- 2.4. The colloidal-crystal-like organization (HEMSLEY, COLLINSON and BRAIN, 1992) can be joint to the previous organization level. But this time its lower or elementary biopolymer units are not well known.
3. The biopolymer units, which can be modelled with fullerenes can be distinguished from the biopolymer systems mentioned above. The relations between the quasi-crystalloid and quasi-equivalent biopolymer systems are under elaboration.

Previously it was believed that the buckyball-like biopolymer units are characteristic for the wall of "peculiar living organisms" such as the genus *Botryococcus*.

Materials and Methods


The pollen grains were collected by Dr. L. TÉCSI from the Botanical Garden of the J. A. University, Szeged, on 20th June 1988. Fresh, non-experimental pollen grains were also used in TEM investigations for comparisons. The experiments started on 20th June 1988 and the different kinds of experiments are the following.

- No 265: 20 mg air dried pollen grains + 1 ml 2-aminoethanol, temperature 30°C , length of time 24^{h} .
- No 280: the pollen grains were heated onto 100°C for 1 hour, after being partially degraded as previously (265).
- No 266: 20 mg air dried pollen grains + 1 ml 2-aminoethanol, temperature 30°C , length of time 24^{h} , washing (H_2O) + 10 ml KMnO_4 aq. dil. 1%, temperature 30°C , length of time 24^{h} .
- No 281: the pollen grains were heated onto 100°C for 1 hour, after being partially degraded as previously (266).
- No 267: 20 mg air dried pollen grains + 1 ml 2-aminoethanol, temperature 30°C , length of time 24^{h} , washing (H_2O) + 10 ml KMnO_4 aq. dil. 1%, temperature 30°C , length of time 48^{h} .


Plate 6.2. ►

Thalictrum flavum L. Recent.

Experiment No: 282, buckyball-like highly organized biopolymer structures from partially degraded and fragmented exine. The superficial electron dense molecular units and the partial degradation process are well illustrated.



- No 282: the pollen grains were heated onto 100 °C for 1 hour, after being partially degraded as previously (267).
- No 268: 20 mg air dried pollen grains + 1 ml 2-aminoethanol, temperature 30 °C, lenght of time 24^h, washing (H₂O) + 10 ml KMnO₄ aq. dil. 1%, temperature 30 °C, lenght of time 24^h, washing (H₂O) + 2 ml acetic acid anhydride, temperature 30 °C, lenght of time 24^h.
- No 283: the pollen grains were heated onto 100 °C for 1 hour, after being partially degraded as previously (268).
- No 269: 20 mg air dried pollen grains + 1 ml 2-aminoethanol, temperature 30 °C, lenght of time 24^h, washing (H₂O) + 10 ml KMnO₄ aq. dil. 1%, temperature 30 °C, lenght of time 24^h, washing (H₂O) + 5 ml methanol, temperature 30 °C, lenght of time 24^h.
- No 284: the pollen grains were heated onto 100 °C for 1 hour, after being partially degraded as previously (289).



Text-fig. 6.1.

Variation statistical graphs of the diameter of the highly organized buckyball-like biopolymer structures. The thick line represents the general average of all measurements, the broken lines represent that of the results by experiments or sometimes by negatives. The numbers of the graphs represent the following: number of experiment, number of the measured biopolymer units, number of the negative.

Ultrathin sections were made on a Porter Blum ultramicrotome with glass knives (post-fixation with OsO_4 aq. dil. embedding in Araldite Durcupan, Fluka). The fragmentation was made with a magnetic stirrer in watered medium, during 30 minutes. The fragmented exines were dropped on a grid covered with collodium pellicle and then they were dried. The electron microscopical investigations were made on a Tesla BS-500 transmission electron microscope, resolution 6 Å.

Results

LM morphology: "16 μ . – Grains cribellate with about 8 pores. Exine with reticulate texture." (ERDTMAN, 1954, p. 120); "6–12 pantoporate" (CLARKE, PUNT and HOEN, 1991, p. 146).

TEM structure of the fresh pollen grains.

Interapertural exine (Plate 6.1., fig. 1). – Tectate, tectum perforated with channels, and ornamented with coni of different size and shape. The surface of the tectum is covered with pollenkitt (cf. HESSE, 1978, p. 20 .. "auf dem Tectum liegen geringfügige Mengen teilweise granulären Pollenkitts...") Infratectal layer columellar. The foot layer is a bit thinner than the tectum. Beneath the foot layer, the interbedded zone was observed (cf. FREAN, 1973, KEDVES and ANTUNOVICS, 1979). The intine is thick, the plasma membrane is thin.

Apertural exine (Plate 6.1., fig. 2). – The ectexine is extremely reduced, and pro parte fragmented (cf. ROLAND, 1966). The intine in this area is thicker than extragerminally; oncus-like.

TRANSMISSION ELECTRON MICROSCOPY OF THE PARTIALLY DEGRADED AND FRAGMENTED EXINES

Experiment No: 265 (Text-fig. 6.1.)

Globular biopolymer units were observed in several kinds of arrangements, linear, network-like, irregular polygons. Diameter: 25 – 35 – 85 Å. The typical buckyball-like biopolymer characteristic features were not observed.


Experiment No: 280

Our TEM data are not sufficient in this respect, this may be the consequence of methodical problems.

Experiment No: 266 (Plate 6.1., fig. 4, text-fig. 6.1., 6.2.)

Globular units of several kinds of arrangements were observed. In highly magnified pictures, not so characteristic buckyball type biopolymer units were observed, large globular units with smaller electron dense superficial molecular systems. In this experiment, the measurements were made on three negatives, the data are not the same;

Negative number	Diameter of the globular units in Å	Diameter of the superficial molecular systems in Å
9751	25 – 40 – 60	5 – 10 – 15
9757	20 – 35 – 60	5 – 10 – 15
9759	20 – 45 – 60	5 – 10 – 15



Experiment No: 281 (Plate 6.1., fig. 3, text-fig. 6.1., 6.2.)

This experiment resulted essentially the previously discussed one; diameter of the globular biopolymer units: 20 – 30 – 60 Å, diameter of the superficial electron dense molecular systems: 5 – 10 – 15 Å.

Experiment No: 267

The qualitative results are similar to those of the experiment No 266, but the biopolymer units were not suitable for quantitative measurements.

Experiment No: 282 (Plate 6.2., text-fig. 6.1., 6.2.)

The buckyball-like biopolymer structures are well shown in our pictures but in several particles of the fragmented exine advanced degradation process was observed. Diameter of the large globular units: 20 – 35, 40 – 60 Å, diameter of the superficial electron dense molecular systems: 5 – 10 – 15 Å.

Experiment No: 268 (Text-fig. 6.1., 6.2.)

Similar to the previously mentioned ones. Diameter of the large globular units: 20 – 35 – 60 Å, diameter of the superficial electron dense molecular systems: 5 – 10 – 15 Å.

Experiment No: 283 (Text-fig. 6.1., 6.2.)

The globular biopolymer units are relatively smaller than in the previous experiments; 15 – 30 – 50 Å, the size of the superficial electron dense molecular systems is constant; 5 – 10 – 15 Å.

Experiment No: 269 (Text-fig. 6.1., 6.2.)

Diameter of the large globular biopolymer units: 25 – 40 – 60 Å. Diameter of the superficial electron dense molecular systems: 5 – 10 – 15 Å.

Experiment No: 284

Advanced disintegration of the biopolymer structure was observed. The quasi-equivalent structures were not in a measurable preservation.

Discussion and Conclusions

1. The buckyball-like biopolymer structure was first observed on *angiosperm* exines. In this way this may occur in other taxa too.
2. The new results underlined newly that the molecular system of the plant cell wall is extremely complicated. Several kinds of structures can be established.
3. The sporopollenin-type plant cell wall is resistant, but as we have pointed it out several times previously this extremely complex molecular system is a dynamically and perpetually altering structure.
4. The chemical compound of the plant cell wall, and its highly organized structures on several organization levels are not unique, either, and it seems that after last time's obtained results, several details need more and more investigations.

◀ Text-fig. 6.2.

Variation statistical graphs of the diameter of the superficial electron dense molecular systems. Thick line represents the general average of all measurements, the broken lines represent of the results by experiment or sometimes by negatives. Numbers of the graphs represent the following: number of experiment, number of the measured electron dense superficial molecular systems, number of the negative.

Acknowledgements

This work was supported by the Grant OTKA 1/3, 104.

References

- BECK, M. és BRAUN, T. (1992): Forradalom a kémiában: a fullerének felfedezése. – Magyar Tudomány 37, 1415–1429.
- CLARKE, G. C. S., PUNT, W. and HOEN, P. P. (1991): The Northwest European Pollen Flora 51 *Ranunculaceae*. – Rev. Palaeobot. Palynol. 69, 117–271.
- ERDTMAN, G. (1954): An Introduction to Pollen Analysis. – Waltham, Mass., U. S. A., Stockholm, Almqvist and Wiksell.
- FREAN, M. L. (1973): Exine stratification and fine structure of the pollen wall of *Croton gratissimus* (BURCH.) subsp. *subgratissimus* (PRAIN) BURTT DAVY. – Pollen et Spores 15, 353–362.
- HEMSLEY, A. R., COLLINSON, M. E. and BRAIN, A. P. R. (1992): Colloidal crystal-like structure of sporopollenin in the megaspore walls of Recent *Selaginella* and similar fossil spores. – Bot. J. of the Linnean Soc. 108, 307–320.
- HESSE, M. (1978): Entwicklungsgeschichte und Ultrastruktur von Pollenkitt und Exine bei nahe verwandten entomophilen und anemophilen *Angiospermensippen*: *Ranunculaceae*, *Hamamelidaceae*, *Platanaceae* und *Fagaceae*. – Pl. Syst. Evol. 130, 13–42.
- HESSE, M. (1985): Hemispheric surface processes of exine and orbicules in *Calluna* (*Ericaceae*). – Grana 24, 93–98.
- KEDVES, M. and ANTUNOVICS, J. (1975): New characteristics in the submicroscopic exine structure of the pollen grains of *Nymphaeaceae* from and evolutionary point of view. – Acta Biol. Szeged. 21, 41–42.
- KEDVES, M., ROJIK, I. and VÉR, A. (1991): Biopolymer organization of the partially degraded oil shale with the fragmentation method. – In: Plant Cell Biology and Development 1 ed.: M. KEDVES, 28–31, Szeged.
- KEDVES, M., ROJIK, I. and VÉR, A. (1992): Ultrastructure and biopolymer organization of the *Botryococcus* colonies from Hungarian alginite. – Workshop on Pyrolysis in Organic Geochemistry, International Workshop, Szeged, Hungary, Abstract, 21, 22.
- KEDVES, M. and TÓTH, A. (1992): Premiers résultats du système de biopolymère stabilisateurs du squelette quasi-cristalloïde de l'exine. – 9. Simposio de Palinologia A. P. L. E., Las Palmas de Gran Canaria – Islas Canarias – 30 Noviembre al 4 Diciembre de 1992, Resúmenes, 20.
- KEDVES, M. and TÓTH, A. (1994): Premiers résultats du système de biopolymère stabilisateurs du squelette quasi-cristalloïde de l'exine. – In: Plant Cell Biology and Development 5 ed.: M. KEDVES, 79–86.
- PENROSE, R. (1979): A class of non-periodic tilings of the plane. – Mat. Int. 2, 32–37.
- ROLAND, F. (1966): Étude de l'ultrastructure des ouvertures: pollens à pores. – Pollen et Spores 8, 409–419.
- ROWLEY, J. R., DAHL, A. O., SENGUPTA, S. and ROWLEY, J. S. (1981): A model of exine substructure based on dissection of pollen and spore exines. – Palynology 5, 107–152.
- SOUTHWORTH, D. (1986): Substructural organization of pollen exines. In: Pollen and Spores: Form and Function. – Linnean Society of London, 61–69.

7. INCOMPLETE AND NON-FIVEFOLD ROTATION OF THE BASIC BIOPOLYMER UNIT OF THE EXINE OF *PINUS GRIFFITHII* MCCLELL

M. KEDVES, A. TÓTH and E. GOTTL

1. Cell Biological and Evolutionary Micropaleontological Laboratory of the Department of Botany of the J. A. University, H-6701, P. O. Box 657, Szeged, Hungary

Abstract

In this paper the results of the incomplete and non-fivefold rotation method applied to the basic pentagonal biopolymer unit of *Pinus griffithii* MCCLELL are presented. This is a new contribution to the method of the basic pentagon biopolymer unit of the sporoderm.

Key words: Palynology, exine, biopolymer structure, rotation method.


Introduction

It was not long after the discovery of SHECHTMAN, BLECH, GRATIAS and CAHN (1984) that in 1988 the presence of the quasi-crystalloid biopolymer structure in living system was first reported from the partially degraded exine of *Pinus griffithii* MCCLELL (KEDVES, 1988). The research program of this subject has split into several directions. One of them is the basic, two-dimensional field which uses the modified MARKHAM rotation method (e. g., KEDVES, 1989, 1990, KEDVES and FARKAS, 1991, etc.).

In the first methodical paper (KEDVES, 1989) the incomplete and non-fivefold rotation of the regular pentagonal polygon biopolymer system in angstrom dimension was also pointed out as a theoretical possibility. Until now, the non-fivefold rotation method has been used in peculiar cases such as the symmetry operations of the TICOS polyhedra modelling to the biopolymer system (KEDVES, 1991) and to the peculiar biopolymer network of the partially degraded and fragmented wall of *Botryococcus braunii* KÜTZ. isolated from the Upper Tertiary oil shale of Hungary (KEDVES, TÓTH and FARKAS, 1993, in print).

Material and Methods

As it has been emphasized several times in a number of publications (cf. SOUTHWORTH, 1986), the basis of partial degradation is the heterogeneity of the different components of the sporoderm, in general, of the ultrastructural elements of the living systems. In this way the solubility and the resistance of the different components is not the same. Using partial degradation before the TEM method, we obtained several



data concerning the biopolymer organization of the plant cell wall on different levels. Since this paper is essentially a new methodical contribution to the actual investigations, we have used the basic regular pentagonal unit, discovered first, which was published in several papers before (cf. KEDVES, 1989).

Results

INCOMPLETE ROTATION METHOD

(Plate 7.1., figs. 1–8, plate 7.2., figs. 1–4)

Designations:

I = Incomplete rotation; the total of the angles of rotation is below 360 °.

P = Primary rotation; the centre of rotation is the middle of one biopolymer unit observed with direct TEM method.

I.P.5.A.5.10/2,1–5. (Plate 7.1., figs. 1,2)

In the case a regular fivefold biopolymer unit rotated with the starting AP axis around the first half of the tenfold rotation (C.P.5.A.5.10). 10/2,1–5 indicate, that the expositions were made at the first five rotation axes as follows:

1st exposition – at the 1st globular biopolymer unit of the regular basic pentagon,

2nd exposition – at the central point of the 1st and 2nd globular biopolymer units of the regular basic pentagon,

3rd exposition – at the 2nd globular biopolymer unit of the regular basic pentagon,

4th exposition – at the central point of the 2nd and 3rd globular biopolymer units of the regular basic pentagon,

5th exposition – at the 3rd globular biopolymer unit of the regular basic pentagon.

Short: this rotation may be designated: A – B – A – B – A. Three globular units and two central points.

I.P.5.A.5.10/2,2–6. (Plate 7.1., figs. 3,4)

1st exposition – at the central point of the 1st and 2nd globular biopolymer units of the regular basic pentagon,

2nd exposition – at the 2nd globular biopolymer unit of the regular basic pentagon,

3rd exposition – at the central point of the 3rd and 4th globular biopolymer units of the regular basic pentagon,

◀ Plate 7.1.

1–8. *Pinus griffithii* McCLELL. Recent. Experiment No: 79, negative no: 7451. Incomplete rotation pictures.

1. I.P.5.A.5.10/2,1–5. 1 Million.

2. I.P.5.A.5.10/2,1–5. 500.000x.

3. I.P.5.A.5.10/2,2–6. 500.000x.

4. I.P.5.A.5.10/2,2–6. 1 Million.

5. I.P.5.A.5.10/2,3–7. 1 Million.

6. I.P.5.A.5.10/2,3–7. 500.000x.

7. I.P.5.A.5.10/2,4–8. 500.000x.

8. I.P.5.A.5.10/2,4–8. 1 Million.

4th exposition – at the 3rd globular biopolymer unit of the regular basic pentagon,
 5th exposition – at the central point of the 3rd and 4th globular biopolymer units of the regular basic pentagon.

Short: this rotation may be designated: B – A – B – A – B. Three central points and two globular biopolymer units.

I.P.5.A.5.10/2,3–7. (Plate 7.1., figs. 5,6)

1st exposition – at 2nd globular biopolymer unit of the regular basic pentagon,

2nd exposition – at the central point of the 2nd and 3rd globular biopolymer units of the regular basic pentagon,

3rd exposition – at the 3rd globular biopolymer unit of the regular basic pentagon,

4th exposition – at the central point of the 3rd and 4th globular biopolymer units of the regular basic pentagon,




Plate 7.2.

- 1–4. *Pinus griffithii* McCLELL. Recent. Experiment No: 79, negative no: 7451. Incomplete rotation pictures.
1. I.P.5.A.5.10/2,5–9. 1 Million.
2. I.P.5.A.5.10/2,5–9. 500.000x.
3. I.P.5.A.5.10/2,6–10. 500.000x.
4. I.P.5.A.5.10/2,6–10. 1 Million.

5th exposition – at the 4th globular biopolymer unit of the regular basic pentagon.
Short: this rotation may be designated: A – B – A – B – A. Three globular units and two central points again.

I.P.5.A.5.10/2,4–8. (Plate 7.1., figs. 7,8)

1st exposition – at the central point of the 2nd and 3rd globular biopolymer units of the regular basic pentagon,

2nd exposition – at the 3rd globular biopolymer unit of the basic regular pentagon,

3rd exposition – at the central point of the 3rd and 4th globular biopolymer units of the regular basic pentagon,

4th exposition – at the 4th globular biopolymer unit of the regular basic pentagon,

5th exposition – at the central point of the 4th and 5th globular biopolymer units of the regular basic pentagon.

Short: this rotation may be designated: B – A – B – A – B. Three central points and two globular units.

I.P.5.A.5.10/2,5–9. (Plate 7.2., figs. 1,2)

1st exposition – at 3rd globular biopolymer unit of the regular basic pentagon,

2nd exposition – at the central point of the 3rd and 4th globular biopolymer units of the basic regular pentagon,

3rd exposition – at the 4th globular biopolymer unit of the regular basic pentagon,

4th exposition – at the central point of the 4th and 5th globular biopolymer units of the basic regular pentagon,

5th exposition – at the 5th globular biopolymer unit of the regular basic pentagon.

Short: this rotation may be designated: A – B – A – B – A. Three globular units and two central points.

I.P.5.A.5.10/2,6–10. (Plate 7.2., figs. 3,4)

1st exposition – at the central point of the 3rd and 4th globular biopolymer units of the basic regular pentagon,

2nd exposition – at the 4th globular biopolymer unit of the regular basic pentagon,

3rd exposition – at the central point of the 4th and 5th globular biopolymer units of the basic regular pentagon,

4th exposition – at the 5th globular biopolymer unit of the regular basic pentagon,

5th exposition – at the central point of the 5th and 1st globular biopolymer units of the basic regular pentagon.

Short: this rotation may be designated: B – A – B – A – B. Three central points and two globular units.


NON-FIVEFOLD ROTATION METHOD OF THE REGULAR BASIC PENTAGONAL BIOPOLYMER UNIT

C.P.5.A.2.2. (Plate 7.3., figs. 1,2)

Interesting secondary points appeared in a peculiar arrangement. This more or less linear biopolymer system is oblique to the PA axis.

C.P.5.A.3.3. (Plate 7.3., figs. 3,4)

In the two circles two times three points of symmetries are reinforced.



C.P.5.A.4.4. (Plate 7.3., figs. 5,6)

Quadrangular points of symmetry appeared with well characterized rotation areas. The discussion of the rotation areas is under preparation in another paper.

C.P.5.A.3.6. (Plate 7.3., fig. 7, plate 7.4., fig. 1)

Interesting circular points of symmetry appeared. These are sexangular in several circles, but the elements of symmetry that appeared in the same circle are not equivalent.

C.P.5.A.7.7. (Plate 7.3., fig. 8, plate 7.4., fig. 2)

This kind of rotation also resulted seven positive and negative points of symmetry together with characteristic rotation areas. But similarly to the previous one, an interesting kind of asymmetry can be observed at the secondary points of symmetry.

C.P.5.A.4.8. (Plate 7.3., fig. 9, plate 7.4., fig. 3)

Negative and positive points of symmetry appeared. The rotation area is almost symmetrical.

C.P.5.A.9.9. (Plate 7.4., figs. 4,5)

This kind of ninefold rotation is essentially the same as the previous one, except for the fact that there are nine positive and nine negative points of symmetry. The rotation area is characteristic.

Secondary rotation picture: C.P.5.A.2.2. – C.S.X_{1/2}.2.2. (Plate 7.4., figs. 6,7)

Several twofold biopolymer units appeared following this kind of secondary rotation. The whole picture and the orientation is similar to those of the primary twofold rotation of the pentagonal biopolymer unit.

Secondary rotation picture: C.P.5.A.2.2. – C.S.X_{1/2}.5.5. (Plate 7.5., figs. 1,4)

Very characteristic fivefold positive and negative points of symmetry appeared after this kind of rotation, together with characteristic rotation areas.

Secondary rotation picture: C.P.5.A.2.2. – C.S.X_{1/2}.4.4. (Plate 7.5., figs. 2,5)

The secondary points that appeared are of tetragonal symmetry with the characteristic rotation areas. The points are more or less duplicate.


Secondary rotation picture: C.P.5.A.4.4. – C.S.X_{2/1}.5.5. (Plate 7.5., figs. 3,6)

Several positive and negative points of symmetry appeared in an unusual, interesting pentagonal arrangement. The delineation of the rotation area is definite.

◀ Plate 7.3

1–9. *Pinus griffithii* MCCLELL. Recent. Experiment No: 79, negative no: 7451. Non-fivefold rotation pictures.

1. C.P.5.A.2.2. 500.000x.
2. C.P.5.A.2.2. 1 Million.
3. C.P.5.A.3.3. 500.000x.
4. C.P.5.A.3.3. 1 Million.
5. C.P.5.A.4.4. 500.000x.
6. C.P.5.A.4.4. 1 Million.
7. C.P.5.A.3.6. 500.000x.
8. C.P.5.A.7.7. 500.000x.
9. C.P.5.A.4.8. 500.000x.



Secondary rotation picture: C.P.5.A.3.6. – C.S.X_{2/1}.3.6. (Plate 7.6., figs. 1,2)

The rotation area is characteristic and sexangular. The positive and negative secondary points of symmetry are generally duplicate.

Secondary rotation picture: C.P.5.A.3.6. – C.S.X_{2/1}.5.5. (Plate 7.6., figs. 3,4)

The secondary points of symmetry are in a pentagonal arrangement. The delineation of the rotation area is not characteristically straight.

Discussion and Conclusions

Regarding the incomplete rotation, our methodical results can be summarized as follows:

	A	B
10/2, 1–5, 7.1.,1,2 A – B – A – B – A	3	2
10/2, 2–6, 7.1.,3,4 B – A – B – A – B	2	3
10/2, 3–7, 7.1.,5,6 A – B – A – B – A	3	2
10/2, 4–8, 7.1.,7,8 B – A – B – A – B	2	3
10/2, 5–9, 7.2.,1,2 A – B – A – B – A	3	2
10/2, 6–10, 7.2.,3,4 B – A – B – A – B	2	3

It is a normal periodicity at this kind of rotation. But when we compare the results of 10/2,1–5, 3–7, 5–9, and 10/2, 2–6, 4–8, 6–10, it is not so easy to distinguish between the similarities of these two groups. It is interesting that at all of the “half tenfold” rotations the results are almost identical with the secondary points of symmetry of a tenfold rotation. The differences can also be established in contrast to the rotation pictures.

The results of the non-fivefold rotation of the basic regular pentagonal biopolymer unit verified again the complexity of the biopolymer system of the sporoderm. To this, the three-dimensional modelling of the quasi-crystalloid biopolymer unit (KEDVES 1991b, 1992, and KEDVES, TÓTH and FARKAS, 1993, in print) contribute useful pieces of information. The arrangement of the units of biopolymer systems seems to have a lot of variations.

◀ Plate 7.4.

- 1–7. *Pinus griffithii* MCCLELL. Recent. Experiment No: 79, negative no: 7451.
- 1–5. Non-fivefold rotation pictures.
 1. C.P.5.A.3.6. 1 Million.
 2. C.P.5.A.7.7. 1 Million.
 3. C.P.5.A.4.8. 1 Million.
 4. C.P.5.A.9.9. 1 Million.
 5. C.P.5.A.9.9. 1 Million.
6. Secondary rotation picture: C.P.5.A.2.2. – C.S.X_{1/2}.2.2. 500.000x.
7. Secondary rotation picture: C.P.5.A.2.2. – C.S.X_{1/2}.2.2. 1 Million.




Plate 7.5.

- 1-6. *Pinus griffithii* McCLELL. Recent. Experiment No: 79, negative no: 7451.
1. Secondary rotation picture: C.P.5.A.2.2. - C.S.X_{1/2}.5.5. 500.000x.
2. Secondary rotation picture: C.P.5.A.4.4. - C.S.X_{2/1}.4.4. 500.000x.
3. Secondary rotation picture: C.P.5.A.4.4. - C.S.X_{2/1}.5.5. 500.000x.
4. Secondary rotation picture: C.P.5.A.2.2. - C.S.X_{1/2}.5.5. 1 Million.
5. Secondary rotation picture: C.P.5.A.4.4. - C.S.X_{2/1}.4.4. 1 Million.
6. Secondary rotation picture: C.P.5.A.4.4. - C.S.X_{2/1}.5.5. 1 Million.




Plate 7.6.

- 1–4. *Pinus griffithii* MCCLELL. Recent. Experiment No: 79, negative no: 7451.
 1. Secondary rotation picture: C.P.5.A.3.6. – C.S.X_{2/1}.3.6. 1 Million.
 2. Secondary rotation picture: C.P.5.A.3.6. – C.S.X_{2/1}.3.6. 500.000x.
 3. Secondary rotation picture: C.P.5.A.3.6. – C.S.X_{2/1}.5.5. 500.000x.
 4. Secondary rotation picture: C.P.5.A.3.6. – C.S.X_{2/1}.5.5. 1 Million.

Acknowledgements

This work was supported by the Grant OTKA 1/3, 104.

References

- KEDVES, M. (1988): Quasi-crystalloid basic molecular structure of the sporoderm. – 7 Internat. Palynol. Congr. Brisbane, Abstracts, 82.
 KEDVES, M. (1989): Méthod d'étude des biopolymères de la paroi pollinique à structure quasi-cristalloïde. – Rev. de Micropaléontologie 32, 226–234.
 KEDVES, M. (1990): Quasi-crystalloid basic molecular structure of the sporoderm. – Rev. Palaeobot. Palynol. 64, 181–186.

- KEDVES, M. (1991a): TICOS polyhedra as a model in the pentasporan organization. In: *Plant Cell Biology and Development* 2, ed.: M. KEDVES, 43–48, Szeged.
- KEDVES, M. (1991b): Three dimensional modelling of the biopolymer structure of the plant cell wall I. In: *Plant Cell Biology and Development* 2, ed.: M. KEDVES, 63–74, Szeged.
- KEDVES, M. (1992): Three dimensional modelling of the biopolymer structure of the plant cell wall II. In: *Plant Cell Biology and Development* 3, ed.: M. KEDVES, 67–87, Szeged.
- KEDVES, M. and FARKAS, E. (1991): Basis of the tertiary rotation and TICOS modelling of the quasi-crystalloid biopolymer skeleton of the plant cell. In: *Plant Cell Biology and Development* 2, ed.: M. KEDVES, 36–42, Szeged.
- KEDVES, M., TÓTH, A. and FARKAS, E. (1993, in print): Experimental investigation of the biopolymer organization of both recent and fossil sporoderms. – Grana,
- SHECHTMAN, D., BLECH, I., GRATIAS, D. and CAHN, J. W. (1984): Metallic phase with long-range order and no translational symmetry. – *Phys. Rev. Lett.* 53, 1951–1953.
- SOUTHWORTH, D. (1986): Substructural organization of pollen exines. In: *Pollen and Spores: Form and Function*, eds.: S. BLACKMORE and I. K. FERGUSON, 61–69, Linnean Soc. London.

8. PREMIERS RÉSULTATS DU SYSTÈME DE BIOPOLYMÈRE STABILISATEUR DU SQUELETTE QUASI-CRISTALLOÏDE DE L'EXINE

M. KEDVES et A. TÓTH

*Laboratoire de Biologie cellulaire et de Micropaléontologie évolutive de l'Institut Botanique de l'Université
J. A. de Szeged, H-6701, B. P. 657, Hongrie*

Sommaire

Le système biopolymère du sporoderme peut être classé en deux composants principaux: 1. Le squelette quasi-cristalloïde qui est un système metastable. 2. Le système des biopolymères stabilisateurs. Ces molécules ou associations des molécules remplissent les cavités et les frustrations – sensu NELSON – du squelette quasi-cristalloïde et empêchent la désintégration spontanée de ce système de biopolymère. Parallèlement avec l'étude du système de biopolymères quasi-cristalloïdes nous avons commencé des recherches pour éliminer le squelette quasi-cristalloïde afin d'obtenir des documents directs par MeT du système stabilisateur. Sur les photos au MeT de l'intine partiellement dégradée des pollens biaillés de *gymnosperme* (*Pinus griffithii* McCLELL.) on a obtenu pour la première fois des informations directes sur le système de biopolymères stabilisateurs de l'unité de PENROSE-I type squelette quasi-cristalloïde. On a établi l'hétérogénéité du système biopolymère des stabilisateurs qui sont selon nos connaissances jusqu'à présent les suivants:

1. Stabilisateur central de forme globulaire au milieu de l'unité centrale (pentagone-dodécaèdre) du squelette de PENROSE-I. Le système du squelette qui entoure le pentagone-dodécaèdre a aussi un stabilisateur central. Mais les relations entre ces deux stabilisateurs centraux ne sont pas encore bien connues.
2. Les frustrations et les autres cavités du squelette quasi-cristalloïde sont remplies par des systèmes de biopolymères, lamellaires ou fibrillaires. Pour mettre en évidence ce système de biopolymère stabilisateur dans les cadres du squelette quasi-cristalloïde nous venons d'élaborer une méthode. Les premiers résultats dans ce sujet présentés dans cette contribution.

Mots clés: Palynologie, actuel, structure biopolymère, stabilisateur.

Introduction

L'étude multidisciplinaire du paroi extrêmement résistant des spores et de grains de pollen a été le sujet depuis longtemps de plusieurs études. Une énorme quantité de publications ont paru avec des conceptions différentes. Vue l'état actuel de nos connaissances dans ce sujet, il nous semble important de distinguer très nettement deux sortes d'études suivant les méthodes fondamentales de la recherche: 1. Les recherches des précurseurs ou en général des composants chimiques, en utilisant les méthodes de la chimie et de la chimico-physique. 2. Le système des biopolymères ou des sous-unités de l'exine, qui peuvent être étudiés par la microscopie électronique à transmission après le processus de la dégradation partielle. Suivant la mesure de la




Planche 8.1.

Pinus griffithii McCLELL.

Détail du système de biopolymère de l'intine partiellement dégradée, par expériment no: 681. No du négative: 430. Plusieurs sortes d'unités de biopolymères sont illustrées. Les globules de diamètre de 6 à 8 Å sont probablement les stabilisateurs centraux d'un pentagone dodécaèdre du PENROSE-I type squelette quasi-cristalloïde. Il y a encore de deux sorte supplémentaires de biopolymères globulaires, les plus petites représentent les "ombres" des atoms. Le système de biopolymères de forme circulaire autour du stabilisateur central remplissent les cavités des "frustrations" concentriques internes. La diamètre de ces circles est de 32,4 à 33,2 Å. Ce système est particulièrement net autour du stabilisateur central "D". Toutes les unités centrales qui ont été utilisées pour des études supplémentaires sont indiquées par des lettres de "A" à "I".

dégradation partielle, on a pu établir des degrés différents dans l'organisation du système de biopolymère, cf. KEDVES (1989a). La base est l'unité de biopolymère de forme de pentagone régulier dans la dimension des angstroms (8–20 Å environ). Ces éléments composent le squelette quasi-cristalloïde qui est le sujet de plusieurs études (cf. GÉVAY et KEDVES, 1989, KEDVES 1989a,b, 1990, 1992, KEDVES, TÓTH et FARKAS, 1992). Ces composants constituent sur un degré supérieur des "sous-unités" comme les structures hélicoïdales ou tubulaires (ABADIE, HIDEUX et ROWLEY, 1986–1987, FLYNN et ROWLEY, 1971, ROWLEY, 1990, ROWLEY, DAHL et ROWLEY, 1981, ROWLEY, DAHL, SENGUPTA et ROWLEY, 1981, ROWLEY, EL-GHAZALY et ROWLEY, 1987), les polygones irréguliers (SOUTHWORTH, 1985a,b, 198a,b) et les éléments globulaires de HESSE (1985). Les structures qui sont similaires aux cristaux colloïdaux publiés par HEMSLEY, COLLINSON et BRAIN (1992) peuvent être classées aussi sur ce niveau.


Du point de vue du sujet de notre étude, les structures quasi-cristalloïdes sont les plus importantes. Suivant les résultats les plus récents, cette organisation peut se rencontrer sur le niveau des molécules aussi (cf. KEDVES, TÓTH et FARKAS, 1992), mais elle n'est jamais connue sur le niveau des sous-unités de l'exine de dimension de nanomètre.

Comme nous l'avons déjà fait remarquer, le squelette de biopolymères quasi-cristalloïdes représente une structure metastable, et par la désintégration de ce système une énergie remarquable peut être libérée. Une complexité des biopolymères remplissent les cavités du squelette quasi-cristalloïde, en assurant la stabilisation relative du paroi sporal. Nous les appelons stabilisateurs. Pour avoir des documents directs de ce système des associations des molécules il est nécessaire de dissoudre le squelette quasi-cristalloïde. Plusieurs essais ont été faits et, en se basant aux résultats de nos expériences jusqu'à maintenant, il nous semble que c'est plus difficile que la dissolution des biopolymères stabilisateurs.

Ce travail va présenter pour la première fois les données directes par la méthode de la microscopie électronique à transmission des stabilisateurs du système quasi-cristalloïde de PENROSE-I type de système biopolymère. Ensuite on a commencé à élaborer une méthode pour l'étude et l'interprétation de l'arrangement des unités stabilisateurs de biopolymères. Il y a lieu de souligner qu'il y a encore beaucoup de choses à faire dans ce sujet, en ajoutant que même le système quasi-cristalloïde n'est pas encore suffisamment connu en tous les détails.

Matière et Méthode

Les grains de pollen de *Pinus griffithii* McCLELL ont été le sujet de ces études, par expérience No 681. La dégradation partielle a été effectuée par diethylether, la méthode en détail a été publiée précédemment, cf. KEDVES et PÁRDUTZ (1992). Comme on a déjà fait remarquer, les négatives ont été prise par un appareil de MeT Opton EM-902 dans le Laboratoire de Microscopie Electronique de l'Institut de Biophysique du Centre de Biologie de l'Académie des Sciences de Hongrie. La négative (No: 430) a été prise au grossissement de 400.000 de l'intine de grain de pollen partiellement dégradée. Il y a lieu de remarquer que sur cette négative on a observé l'image des molécules, et parmi celles-ci une de forme de pentagone a été le sujet des opérations de symétrie, cf. KEDVES, TÓTH et FARKAS (1992).



Résultats

Pour avoir une aspect général, on a préparé des photos au grossissement de 5 Million de la négative entière. Sur cette grande photo laquelle a été composée de plusieurs parties on a pu observer très clairement des unités globulaires de plusieurs dimensions. Pour la publication nous avons utilisé des photos à 1 Million (Planche 8.1.). Certaines, mais en particulier une ("D") est entourée d'un système de biopolymère de forme circulaire. Un système biopolymère similaire, mais pas tellement complet a été encore observé ("B, F, G, H"). Nous avons pensé que les unités globulaires grandes, de taille de 6 à 8 Å, sont les unités stabilisatrices centrales, qui se placent au milieu du pentagon-dodécaèdre central d'un PENROSE-I type squelette de biopolymère.

Pour étudier la disposition de ces unités de biopolymères dans le système entière, comme premier essai pour la méthode de recherche nous avons effectué les démarches ci-dessous:

1. En tenant compte des unités de biopolymères représentées sur la Planche 8.1., et les schémas (Fig. 8.1. et 8.2.), ensuite en se basant sur nos résultats antérieurs concernant le squelette quasi-cristalloïde en deux mais en particulier en trois dimensions, nous avons pris pour base la photo prise du modèle de PENROSE-I type quasi-cristalloïde squelette à trois dimensions. Comme nous l'avons déjà fait remarquer, nos modèles à trois dimensions ont été préparés dans l'échelle des unités de biopolymères observées sur les premières photos de MeT. De la photo, prise en face du modèle à trois dimensions de PENROSE-I type squelette, nous avons préparé des schémas dans la mesure de 5 et de 1 million. Voir ce dernier sur la figure 8.1. et 8.2.
2. Ce schéma pensons-nous représente le squelette quasi-cristalloïde de PENROSE-I type unité en deux dimensions.
3. Ensuite nous avons placé ce schéma au milieu d'un biopolymère globulaire (qui est selon notre hypothèse le stabilisateur central du PENROSE-I unité), et ensuite tourné jusqu'à la position où le maximum des unités voisines soit en symétrie avec le schéma du squelette (Fig. 8.1. et 8.2.).

◀ Fig. 8.1.

Pinus griffithii MCCLELL.

Schéma préparé de la photo de la planche 8.1., indiquant l'emplacement du système biopolymère des stabilisateurs centraux. Un autre type de schéma qui a été préparé d'une photo pris d'un modèle à trois dimensions d'une unité de type PENROSE-I squelette dans la même échelle de la photo; au grossissement d'1 Million Plusieurs sortes de connexions du squelette quasi-cristalloïde avec les stabilisateurs sont représentées.

Connexion de type I.: Exemple: Biopolymères "A" et "B". C'est la connexion la plus enchainée. Les stabilisateurs centraux des unités connectées du PENROSE-I type squelette se placent sur les systèmes de biopolymères circulaires. Voir encore deux unités globulaires autour du biopolymère "D".

Connexion de type II.: Exemple: Biopolymères "G", "H" et "I". Dans ce cas-là, le contour extérieur du squelette de type PENROSE-I touche le centre (le pentagone ou le stabilisateur central) du squelette voisin.

Connexion de type III.: Exemple: Biopolymère "B" et "D". Les contours extérieures sont en connexion de deux squelettes de PENROSE-I type voisin.




Fig. 8.2.

Pinus griffithii MCCLELL.

Schéma préparé de la photo de la planche 8.1., indiquant tous les points de symétrie importants. Les squelettes de type PENROSE-I des biopolymères stabilisateurs centraux "A", "C", "D" et "E" sont indiqués. Il est à remarquer que suivant ces schémas les stabilisateurs circulaires sont en connexion.

4. Suivant cette méthode, nous avons pu distinguer pour le moment trois positions principales des unités de PENROSE-I type squelette y compris leur stabilisateur central.

4.1. Connection de type I (Fig. 8.1. "A" et "B")

C'est la connection la plus proche entre deux unités. Il est à remarquer que les pentagones centraux avec les unités "A" et "B" sont en position particulière. Un sommet de tous les deux pentagones atteint le stabilisateur central voisin. En analysant cette partie de schéma ("A" et "B", fig. 8.1.) on peut établir encore des symétries intéressantes supplémentaires.

4.2. Connection de type II (Fig. 8.1., "G", "H" et "I", et fig. 8.2., "A", "C", "D" et "E")

Ces connections dans les cas typiques peuvent être caractérisées avec les traits ci-dessous:

Les systèmes de biopolymères stabilisateurs circulaires voisins sont en connexion tangentielle (Fig. 8.2.).

D'autre part, le contour avec un losange périphérique du schéma du squelette PENROSE-I touche le stabilisateur voisin.

Il y a deux sortes de conditions pour ce type de connexion:

1. Une distance convenable entre les deux stabilisateurs centraux; $A - C = 29 \text{ \AA}$, $C - D = 33 \text{ \AA}$, $D - E = 31 \text{ \AA}$, $G - H = 28 \text{ \AA}$, $H - I = 32 \text{ \AA}$.

2. L'angle de trois stabilisateurs voisins semblait aussi important; $AC - CD = 144^\circ$, $CD - DE = 130^\circ$, $GH - HI = 145^\circ$.

Nous avons trouvé une connexion pas typique, voir D et F, fig. 8.1. Les systèmes de biopolymères stabilisateurs circulaires ne se touchent pas, mais les angles extérieurs du schéma sont en connexion avec le pentagone central voisin.

4.3. Connection de type III (Fig. 8.1., "B" et "D")

Dans ce cas-là les contours extérieurs sont en connection tangentielle.

5. Pour avoir des aspects plus amples concernant la complexité du système biopolymère du paroi pollinique, nous avons mis en évidence certains éléments de la symétrie, en particulier sur la fig. 8.1.

Discussion et Conclusions

1. Comme nous avons déjà fait remarquer c'est le premier document direct concernant le système de biopolymère stabilisateur du squelette quasi-cristalloïde. Suivant nos résultats précédents sur le squelette quasi-cristalloïde, il est évident que le système stabilisateur est beaucoup plus complexe que nous n'avons pu le mettre en évidence jusqu'ici. En effet nous avons des informations de deux sortes de stabilisateurs seulement.

2. La méthode présentée ici est aussi pour une sorte de système biopolymère. Par exemple pour les éléments lamellaires ou en particulier le système biopolymère de la surface il faut élaborer une autre méthode.

Remerciements

Le texte français a été corrigé par Mme. Zs. GÉCSEG, nous venons lui exprimer nos remerciements les plus sincères pour son aide cordiale. Ces recherches ont été effectuées avec le concours du OTKA 1/3 - 104.

Bibliographie

- ABADIE, M., HIDEUX, M. and ROWLEY, J. R. (1986–1987): Ultrastructural cytology of the anther. II. Proposal for a model of exine considering a dynamic connection between cytoskeleton, glycolemma and sporopollenin – synthesis. – *Ann. Sci. Nat. Bot. Paris* 8, 1–16.
- FLYNN, J. J. and ROWLEY, J. R. (1971): Wall microtubules in pollen grains. – *Zeiss Inform.* 76, 40–45.
- GÉVAY, G. and KEDVES, M. (1989): A structural model of the sporopollenin based on dodecahedrane units. – *Acta Biol. Szeged.* 35, 53–57.
- HEMSLEY, A. R., COLLINSON, M. E. and BRAIN, A. P. R. (1992): Colloidal crystal-like structures of sporopollenin in the megaspore walls of Recent *Selaginella* and fossil spores. – *Bot. J. Linnean Soc.* 108, 307–320.
- HESSE, M. (1985): Hemispheric surface processes of exine and orbicules in *Calluna* (*Ericaceae*). – *Grana* 24, 93–98.
- KEDVES, M. (1989a): Quasi-crystalloid biopolymer structures of the sporoderm and its highly organized degrees. – *Acta Biol. Szeged.* 35, 59–70.
- KEDVES, M. (1989b): Méthode d'étude des biopolymères de la paroi pollinique à structure quasi-cristalloïde. A method of investigation of the quasi-crystalloid structure of the pollen wall biopolymers. – *Rev. de Micropaléont.* 32, 226–234.
- KEDVES, M. (1990): Quasi-crystalloid basic molecular structure of the sporoderm. – *Rev. Palaeobot. Palynol.* 64, 181–186.
- KEDVES, M. (1992): Three dimensional modelling of the biopolymer structure of the plant cell wall. II. In: *Plant Cell Biology and Development* 3, ed.: M. KEDVES, 67–87, Szeged.
- KEDVES, M. and PÁRDUTZ, Á. (1992): Transmission electron microscopy of partially dissolved exines of different bisaccate *gymnosperm* pollen grains. In: *Plant Cell Biology and Development* 3, ed.: M. KEDVES, 38–66, Szeged.
- KEDVES, M., TÓTH, A. and FARKAS, E. (1992): Experimental investigation of the biopolymer organization of the sporoderm (recent and fossil). – 8th I. P. C. Aix-en-Provence, Program and Abstracts, 75.
- NELSON, D. R. (1986): Quasicrystals. – *Scientific American* 254, 43–51.
- ROWLEY, J. R. (1990): The fundamental structure of the pollen exine. – *Pl. Syst. Evol.* (Suppl. 5), 13–29.
- ROWLEY, J. R., DAHL, A. O. and ROWLEY, J. S. (1981): Substructure in exines of *Artemisia vulgaris* (*Asteraceae*). – *Rev. Palaeobot., Palynol.* 25, 1–38.
- ROWLEY, J. R., DAHL, A. O., SENGUPTA, A. S. and ROWLEY, J. S. (1981): A model of exine substructure based on dissection of pollen and spore exines. – *Palynology* 5, 107–152.
- ROWLEY, J. R., EL-GHAZALY, G. and ROWLEY, J. A. (1987): Microchannels in the pollen grain exine. – *Palynology* 11, 1–21.
- SOUTHWORTH, D. (1985a): Pollen exine substructure. I. *Lilium longiflorum*. – *Amer. J. Bot.* 72, 1274–1283.
- SOUTHWORTH, D. (1985b): Pollen exine substructure. II. *Fagus sylvatica*. – *Grana* 24, 161–166.
- SOUTHWORTH, D. (1986a): Pollen exine substructure. III. *Juniperus communis*. – *Can. J. Bot.* 64, 983–987.
- SOUTHWORTH, D. (1986b): Substructural organization of pollen exines. In: *Pollen and Spores: Form and Function*, 61–69.

9. LIST OF PUBLICATION OF M. KEDVES II.

compiled by

E. GOTTL

Cell Biological and Evolutionary Micropaleontological Laboratory of the Department of Botany of the J. A. University, H-6701, P. O. Box 657, Szeged, Hungary

1975

--: Sur les problèmes de la structure et de la nomenclature de l'exine des pollens des *Angiospermes* fossiles. – Soc. bot. Fr., Coll. Palynologie 122, 69–73.

Key words: Palynology, fossil, *Angiospermatophyta*, exine stratification, nomenclature.

KEDVES, M.–ANTUNOVICS, J.: New characteristics in the submicroscopic exine structure of the pollen grains of *Nymphaeaceae* from an evolutionary point of view. – Acta Biol. Szeged. 21, 41–42.

Key words: Palynology, recent, *Angiospermatophyta*, exine ultrastructure.

KEDVES, M.–HEGEDŰS, M.: Pollen grains of the *Interporopollenites* fgen. from sediments of the Upper Cretaceous period in Portugal. – Acta Biol. Szeged. 21, 43–62.

Key words: Palynology, fossil, n. fssp., Cretaceous, Portugal.

KEDVES, M.–RADVÁNSZKI, M.: The application of scanning electron microscopical method in some plant microfossils. – Acta Bot. Acad. Sci. Hung. 21, 51–59.

Key words: Palynology, fossil, SEM.

STANLEY, E. A.–KEDVES, M.: Electronmicroscopical investigations of the *Normapolles* group and some other selected European and North American *Angiosperm* pollen I. – Pollen et Spores 17, 233–271.

Key words: Palynology, fossil, *Angiospermatophyta*, TEM, SEM, Cretaceous, Portugal, Eocene, France, Hungary, USA.

1976

--: Scanning electron-microscopic investigations on the pollen grains of the Operculati VENK. et GÓCZ. 1964. – Acta Biol. Szeged. 22, 29–36.

Key words: Palynology, fossil, *Gymnospermatophyta*, SEM, Triassic, Jurassic, Hungary, Cretaceous, Portugal.

KEDVES, M.–STANLEY, E. A. a: Electronmicroscopical investigations of the *Normapolles* group and some other selected European and North American *Angiosperm* pollen II. – Pollen et Spores 18, 105–127.

Key words: Palynology, fossil, *Angiospermatophyta*, TEM, SEM, Eocene, France, Hungary, USA.

KEDVES, M.–STANLEY, E. A. b: Electron-microscope investigations of the form-genus *Pentapollenites* KRUTZSCH 1958, and its re-establishment as a valid genus. – Pollen et Spores 18, 289–297.

Key words: Palynology, fossil, *Angiospermatophyta*, TEM, SEM, Eocene, Hungary.

Book reviews

- a: KRUTZSCH, W.: Atlas der mittel- und jungtertiären dispersen Sporen- und Pollen- sowie der Mikroplanktonformen des nördlichen Mitteleuropas. Lieferung VI. Coniferenpollen (*Saccites* und "*Inaper-turates*"). -- VEB G. Fischer Verlag, Jena, 1971. -- Bot. Közlem. 63, 41.
- b: PANKOW, H.: Algenflora der Ostsee. II. Plankton. -- VEB G. Fischer, Jena, 1976. -- Bot. Közlem. 63, 234.
- c: RHEINHEIMER, G.: Mikrobiologie der Gewässer. -- VEB G. Fischer, Jena, 1975. -- Bot. Közlem. 63, 234.

1977

- a: Contribution de l'ornementation en stries concentriques à la connaissance des microfossiles. -- Pollen et Spores 19, 404-414.
Key words: Palynology, fossil, SEM, Upper Tertiary, Angola.
- b: Electronmicroscopical examinations of fossil *Angiospermatophyta* pollen grains from the Paleocene and the Middle Eocene. -- Acta Bot. Acad. Sci. Hung. 23, 97-103.
Key words: Palynology, fossil, TEM, SEM, Paleocene, France, Eocene, Hungary.

1978

- a: Paleogene fossil sporomorphs of the Bakony Mountains III. -- Studia Biologica Academiae Scientiarum Hungaricae 15, Akadémiai Kiadó, Bp.
Key words: Palynology, fossil, n.fgen., n.fssp., Paleogene, Hungary.
- b: Ultrastructure investigations into fossil *Salviniaceae* spores. -- Acta Biol. Szeged. 24, 19-22.
Key words: Palynology, fossil, *Pteridophyta*, ultrastructure, Cretaceous, Egypt.
- c: Palynological investigations into sediments of the Lower Paleogene period in Bulgaria. -- Acta Biol. Szeged. 24, 23-30.
Key words: Palynology, fossil, Paleogene, Bulgaria.
- d: On nomenclature problems of the Prequarternary fossil sporomorphs. -- IV. Int. Palynol. Conf., Lucknow (1976-77) 1, 191-193.
Key words: Palynology, fossil, exine structure and nomenclature.
- KEDVES, M.-SIMONCSICS, P.: The sporomorphs of an Angolan brown coal. -- Acta Bot. Acad. Sci. Hung. 24, 69-89.
Key words: Palynology, fossil, n. fssp., SEM, Upper Tertiary, Angola.

1979

- a: Scanning electron microscopy of some selected recent *Amentiflorae* pollens I. -- Acta Bot. Acad. Sci. Hung. 25, 75-82.
Key words: Palynology, recent, *Amentiflorae*, SEM.
- b: Données stratigraphiques sur les *Angiospermes* du Crétacé supérieur d'Europe. -- Paleobiologie continentale 10, 18-22.
Key words: Palynology, fossil, stratigraphy, Upper Cretaceous, Europe.
- c: Intraspecific morphological variations at recent *Angiospermatophyta* pollen grains. -- Acta Biol. Szeged. 25, 65-68.
Key words: Palynology, recent, *Angiospermatophyta*, intraspecific morphological variations.
- d: Scanning electron-microscopical investigations into the sporomorphs of the coal layers in the Dorog Basin. -- Acta Biol. Szeged. 25, 35-44.
Key words: Palynology, fossil, SEM, Eocene, Hungary.
- e: Testing of the spores in the *Equisetum* genus. (In Hungarian, summary in English). -- Bot. Közlem. 66, 195-203.
Key words: Palynology, recent, *Equisetum*, LM, SEM.

-- f: Palynological investigations on sediments of the Lower Danian (Fish Clay, Denmark) I. – *Acta Miner.-Petr. Szeged* 24, 167–186.

Key words: Palynology, fossil, *Angiospermatophyta*, n. fgen., n. fspp., Cretaceous – Tertiary, Denmark.

KEDVES, M.–DINIZ, F. a: Les pollens d'*Angiospermes* du Crétacé de Vila Flor, Portugal. Genres de forme *Atlantipollis* et *Limaipollenites*. – *Boletim da Sociedade Geológica de Portugal* 21, 203–216.

Key words: Palynology, fossil, *Angiospermatophyta*, n. fspp., Upper Cretaceous, Portugal.

KEDVES, M.–DINIZ, F. b: Étude au microscope électronique à balayage de quelques espèces du genre de forme *Interporopollenites* du Crétacé d'Arada, Portugal. – *Boletim da Sociedade Geológica de Portugal* 21, 217–226.

Key words: Palynology, fossil, SEM, Upper Cretaceous, Portugal.

KEDVES, M.–PITTAU, P.: Contribution à la connaissance des pollens des *Normapollis* du "groupe papilloïde" du Crétacé supérieur du Portugal. – *Pollen et Spores* 21, 169–209.

Key words: Palynology, fossil, n.fgen., n. fspp., Upper Cretaceous, Portugal.

Book reviews

-- a: Ettl, H., Gerloff, J.–Heynig, H.: Süßwasserflora von Mitteleuropa 3, Ettl, H.: *Xanthophyceae* 1. Teil. – VEB Gustav Fisher Verlag, Jena, 1978. – *Bot. Közlem.* 66, 204.

-- b: Nilsson, S., Pragowski, J.–Nilsson, L.: Atlas of Airborne Pollen Grains and Spores in Northern Europe. – Natur och Kultur Stockholm, 1977. – *Bot. Közlem.* 66, 204.

1980

-- a: Morphological investigation on recent *Palmae* pollen grains. – *Acta Bot. Acad. Sci. Hung.* 26, 339–373.

Key words: Palynology, recent, *Palmae*, LM, SEM.

-- b: Palynological investigations on Austrian Upper Cretaceous and Lower Tertiary sediments. – *Acta Biol. Szeged* 26, 63–77.

Key words: Palynology, fossil, n.fgen., n. fspp., Upper Cretaceous – Eocene, Austria.

-- c: Evolutionary problems of Early *Brevaxones* pollen genera. – Internat. Palynol. Conf., Abstracts, 198.

Key words: Palynology, fossil, *Angiospermatophyta*, evolution.

-- d: Palynological investigations on sediments of the Lower Danian (Fish Clay, Denmark) II. – *Acta Miner.-Petr.* 24, 355–376.

Key words: Palynology, fossil, *Pteridophyta*, *Gymnospermatophyta*, n. fgen., n. fspp., Cretaceous – Tertiary, Denmark.

-- e: Les pollens du genre de forme *Complexipollis* W. KR. 1959 em. TSCHUDY 1973 du Cenomanien supérieur de Vila Flor (Portugal). – *Revista Española de Micropaleontología* 12, 469–488.

Key words: Palynology, fossil, n. fspp., Upper Cretaceous, Portugal.

KEDVES, M.–HERNGREEN, G. F. W.: Palynology of the stratotype of the Maestrichtian and the Gulpen Formation, ENCI Section, Maastricht, The Netherlands. – *Pollen et Spores* 22, 483–544.

Key words: Palynology, fossil, n. fgen., n. fspp., Maastrichtian, The Netherlands.

1980–1981

KEDVES, M.–DINIZ, F.: Contribution à la connaissance des pollens d'*Angiospermes* du Crétacé supérieur du Portugal. – *Bol. Soc. Geol. Portugal* 22, 19–39.

Key words: Palynology, fossil, n.fgen., n. fspp., Upper Cretaceous, Portugal.

1981

-- a: The evolutionary significance of the *angiospermous* exine ultrastructure and sculpture. – Inter. Symp. Concept. Meth. Paleo. Barcelona, 75–83.

Key words: Palynology, fossil, *Angiospermatophyta*, TEM, SEM, evolution.

-- b: Études palynologiques sur les sédiments préquaternaires de l'Égypte. Néogène I. – Grana 20, 119–130.

Key words: Palynology, fossil, n. fspp., Neogene, Egypt.

-- c: Definitions of, evolutionary trends within, and classification of early *brevaxonate* pollen. – Rev. Palaeobot. Palynol. 35, 149–154.

Key words: Palynology, fossil, *Angiospermatophyta*, evolution.

-- d: Letter of Hungarian Palynologist. – Japanese Journal of Palynology 27, 70.

Key words: Palynology, fossil, *Probrevaxones*, Upper Cretaceous, Portugal.

-- e: Scanning electron-microscopic investigations on the sporomorphs of the Upper Pannonian in Hungary. – Acta Biol. Szeged. 27, 89–103.

Key words: Palynology, fossil, SEM, Upper Pannonian, Hungary.

KEDVES, M.–DINIZ, F.: *Probrevaxones* a new pollen group for the first *Brevaxones* form-genera from the Upper Cenomanian of Portugal. – Acta Bot. Acad. Sci. Hung. 27, 383–402.

Key words: Palynology, fossil *Brevaxones*, n. fgen., n. fspp., Upper Cretaceous, Portugal.

KEDVES, M.–PÁRDUTZ, Á. a: Transmission electron microscopic (TEM) investigations on Upper Cretaceous spores from Vila Flor (Portugal). – Acta Biol. Szeged. 27, 105–115.

Key words: Palynology, fossil, *Pteridophyta*, TEM, Upper Cretaceous, Portugal.

KEDVES, M.–PÁRDUTZ, Á. b: Études au microscope électronique à transmission des exines des premiers *Brevaxones*. – Revista Española de Micropaleontología 13, 273–288.

Key words: Palynology, fossil, *Angiospermatophyta*, TEM, Upper Cretaceous, Portugal.

1982

-- a: History of the paleophytogeographical regions based on plant microfossils. – Japanese Journal of Palynology 28, 22.

Key words: Palynology, fossil, *Angiospermatophyta*, Paleophytogeography.

-- b: Historia de las regiones paleofitogeográficas a partir de los datos palinológicos. – IV. Simposio de Palinología, Programa, Resúmenes y liste de participantes, 23.

Key words: Palynology, fossil, Paleophytogeography, evolution.

-- c: Studies on the pollen grains of recent *Castaneoideae*. I. – Acta Biol. Szeged. 28, 1–4.

Key words: Palynology, recent, *Castaneoideae*.

KEDVES, M. J.–PÁRDUTZ, Á. a: Ultrastructural investigations of the Early *Normapolles* taxa *Complexiopollis* and *Limaipollenites*. – Palynology 6, 149–159.

Key words: Palynology, fossil, *Angiospermatophyta*, Upper Cretaceous, Portugal.

KEDVES, M.–PÁRDUTZ, Á. b: Complex studies on the pollen grains of *Elaeagnus angustifolia* L. – Acta Biol. Szeged. 28, 1–4.

Key words: Palynology, recent, *Elaeagnus*, LM, TEM, SEM.

KEDVES, M.–RUSSELL, D. E.: Palynology of the Thanetian layers of Menat. – Palaeontographica B, 182, 87–150.

Key words: Palynology, fossil, n. fgen., n. fspp., Paleocene, Menat, France.

Book review

--: CASPER, S. J.–KRAUSCH, D.: *Pteridophyta* and *Anthophyta* 2. Teil: *Saururaceae* bis *Asteraceae*. In: Süßwasserflora von Mitteleuropa 24. (eds.: Ettl, H.–Gerloff, J.–Heynig, H.). – VEB Gustav Fischer Verlag, Jena, 1981. – Bot. Közlem. 69, 14.

1983

-- a: La stratification de l'exine et la morphologie des *Normapolles*. – Physio-Géo 6, 53–67.

Key words: Palynology, fossil, *Normapolles*, *Angiospermatophyta*, exine stratification, morphology.

-- b: Études palynologiques sur les sédiments préquaternaires de l'Égypte. Néogène II. – Grana 22, 39–49.

Key words: Palynology, fossil, n. fgen., n. fspp., Neogene, Egypt.

-- c: L'histoire des régions paléophytogéographiques d'après les données palynologiques. In: Actas del IV Simposio de Palynologia APLE - Barcelona, 7-9 Octobre 1982, eds.: SOLÉ DE PORTA, N.-SUÁREZ CERVERA, M., 337-349.

Key words: Palynology, fossil, Paleophytogeography, evolution.

-- d: Étude paléobotanique sur les schistes pétrolifères du Tertiaire Supérieur de Hongrie. - Rev. de Micropaléontologie 26, 48-53.

Key words: Palynology, fossil, *Botryococcus*, oil shale, Neogene, Hungary.

-- e: The phylogenetic and taxonomic questions of the pollen grains of *Angiospermatophyta*. (In Hungarian, summary in English). - Bot. Közlem. 70, 13-17.

Key words: Palynology, fossil, *Angiospermatophyta*, evolution, Taxonomy.

-- f: Beszámoló az A.P.L.E. IV. Palinológiai Szimpóziumáról (Barcelona, 1982. október 7-9.). - Bot. Közlem. 70, 115-116.

Key words: Palynology, APLE Symposium, review.

-- g: A paleofitogeográfiai régiók fejlődéstörténete a növényi mikrofosztiliák alapján. - MTA Biol. Oszt. Köz. 25, 697-704.

Key words: Palynology, fossil, Paleophytogeography, evolution.

-- h: Development of the European *Brevaxones* pollen grains and the main stages of their evolution during the Lower and Middle Senonian. - Pollen et Spores 25, 487-597.

Key words: Palynology, fossil, *Brevaxones*, evolution, Upper Cretaceous, Europe.

-- i: *Endoinfundibulapollis distinctus* R. TSCHUDY 1975, from the Upper Cretaceous from the southern part of Hungary. First occurrence of this form-genus from Europe. - Acta Biol. Szeged. 29, 199-200.

Key words: Palynology, fossil, Upper Cretaceous, Hungary.

KEDVES, M.-DINIZ, F.: Les *Normapollis* du Crétacé supérieur en Europe: Implications Paléobiogéographiques. - Geobios 16, 329-345.

Key words: Palynology, fossil, n. fgen., n. fspp., Paleophytogeography, Upper Cretaceous, Europe.

KEDVES, M.-PÁRDUTZ, Á. a: Scanning electron microscopy of some selected recent *Amentiflorae* pollens II. - Acta Biol. Szeged. 29, 67-76.

Key words: Palynology, recent, *Amentiflorae*, SEM.

KEDVES, M.-PÁRDUTZ, Á. b: Studies on the pollen grains of recent *Castaneoideae* II. - Acta Biol. Szeged. 29, 77-88.

Key words: Palynology, recent, *Castaneoideae*, TEM, SEM.

KEDVES, M.-PÁRDUTZ, Á. c: Electron microscope investigation of the Early *Normapollis* pollen genus *Atlantopollis*. - Palynology 7, 153-169.

Key words: Palynology, fossil, *Atlantopollis*, TEM, SEM, Upper Cretaceous, Portugal.

Book review

--: BRAUNE, W., LEMAN, A.-TAUBERT, H.: Pflanzenanatomisches Praktikum II. - VEB Gustav Fisher Verlag, Jena, 1982. - Bot. Közlem. 70, 12.

1984

-- a: Upper Cretaceous sporomorphs from the southern part of Hungary (Csávoly). - Acta Biol. Szeged. 30, 75-89.

Key words: Palynology, fossil, n. fgen., n. fspp., Upper Cretaceous, Hungary.

-- b: Étude palynologique d'un lignite tertiaire de Blão, Viet-nam -I-. - Acta Biol. Szeged. 30, 91-105.

Key words: Palynology, fossil, n. fgen., n. fspp., Neogene, Viet-nam.

-- c: Études palynologiques sur les sédiments préquaternaires de l'Égypte. Danien. - Revista Española de Micropaleontología 16, 43-50.

Key words: Palynology, fossil, Upper Cretaceous, Danien, Egypt.

-- d: Ultrastructure de la paroi des spores des *Lycopodiaceae* du Crétacé supérieur d'Égypte. - Revue de Micropaléontologie 27, 189-195.

Key words: Palynology, fossil, *Lycopodiaceae*, TEM, Upper Cretaceous, Egypt.

-- e: Cretaceous sporomorphs from Gubbio, Italy. – *Palaeontographia Italica* 73, 34–40.

Key words: Palynology, fossil, n. fsp., Cretaceous, Italy.

KEDVES, M.–KÖRMÖCZI, L.: Los problemas de la preservación de esporomorfos bajo condiciones diferentes. – V Simposio de Palinología APLE, Córdoba 1984, Resúmenes, 25.

Key words: Palynology, Holocene, spore-pollen preservation.

KEDVES, M., SOLÉ DE PORTA, N., DE PORTA, J.–CIVIS, J.: Estudio de los sedimentos del Barranco de La Posa (Lérida, España). – V Simposio de Palinología APLE, Córdoba 1984, Resúmenes, 26.

Key words: Palynology, fossil, Upper Cretaceous, Spain.

1985

-- a: Structural modification of degraded fossil sporomorphs. – *Micropaleontology* 31, 175–180.

Key words: Palynology, recent, fossil, TEM partial degradation.

-- b: Étude palynologique d'un lignite tertiaire de Blău, Viet-nam –II–. – *Acta Biol. Szeged.* 31, 97–113.

Key words: Palynology, fossil, n. fgen., n. fssp., Neogene, Viet-nam.

-- c: The present day state of Upper Cretaceous palaeophytogeography on palynological evidence. – *Acta Biol. Szeged.* 31, 115–127.

Key words: Palynology, fossil, Paleophytogeography, Upper Senonian.

-- d: LM, TEM and SEM investigations on recent inaperturate *Gymnospermatophyta* pollen grains. – *Acta Biol. Szeged.* 31, 129–146.

Key words: Palynology, recent, *Taxodiaceae*, *Cupressaceae*, LM, TEM, SEM.

-- e: Études palynologiques sur les sédiments préquaternaires de l'Égypte. Oligocène. – *Revista Española de Micropaleontología* 17, 333–346.

Key words: Palynology, fossil, Oligocene, Egypt.

KEDVES, M.–KÖRMÖCZI, L.: Sur les problèmes de conservation des sporomorphes dans des conditions différentes. – *An. Asoc. Palinol. Leng. Esp.* 2, 263–271.

Key words: Palynology, Holocene, Hungary.

KEDVES, M., SOLÉ DE PORTA, N., DE PORTA, J.–CIVIS, J.: Estudio palinológico de los sedimentos Maastrichtienses del Barranco de La Posa (Prepirineo, Lérida, España). – *An. Asoc. Palinol. Leng. Esp.* 2, 247–253.

Key words: Palynology, fossil, Upper Cretaceous, Spain.

KEDVES, M.–SZEDERKÉNYI, T.: The importance of the spore-pollen investigations in the recognition of the radioactive element content of the lake mud. – *Acta Biol. Szeged.* 31, 215–216.

Key words: Organic microfossils, radioactive materials.

PORTA, DE J., KEDVES, M., SOLÉ DE PORTA, N.–CIVIS, J.: Palinología del Maastrichtiense del Barranco de La Posa (Lérida, España). Problemática regional. – *Rev. Inv. Geol.* 40, 5–28.

Key words: Palynology, fossil, n. fssp., Upper Cretaceous, Spain.

1986

-- a: Introduction to the Palynology of pre-Quaternary deposits. Part I. – *Studia Biologica Academiae Scientiarum Hungaricae* 19, Akadémiai Kiadó, Budapest.

Key words: Palynology, general establishments.

-- b: Introduction to the Palynology of pre-Quaternary deposits Part II. – *Studia Biologica Academiae Scientiarum Hungaricae* 20, Akadémiai Kiadó, Budapest.

Key words: Palynology, fossil, Precambrian – Upper Tertiary.

-- c: Paleogene fossil sporomorphs of the Bakony Mountains IV. – *Studia Biologica Academiae Scientiarum Hungaricae* 21, Akadémiai Kiadó, Budapest.

Key words: Palynology, fossil, stratigraphy, Paleogene, Hungary.

-- d: Aspects and problems of the examination of the fossil spores and pollen grains. – IV. Magyar Növényanatómiai Szimpózium, Abstract, 15.

Key words: Palynology, fossil, aspects, problems.

-- e: Palynological investigations of prequaternary sediments of Egypt. Lower part of the Nubian Sandstone in the Kharga Oasis. – *Z. geol. Wiss. Berlin* 14, 331–355.

Key words: Palynology, fossil, n. fgen., n. fspp., LM, TEM, Jurassic, Egypt.

-- f: Études palynologiques sur les sédiments préquaternaires de l'Égypte. Eocène. – Revista Española de Micropaleontologia 18, 5–26.

Key words: Palynology, fossil, n. fgen., n. fspp., Eocene, Egypt.

-- g: Dégénération expérimentale de la paroi pollinique. – VI. Simp. de Palinol. APLE Resúmenes, 20.

Key words: Palynology, fossil, biopolymer structure.

-- h: Dégénération expérimentale des colonies du genre *Botryococcus* des schistes pétrolifères du Tertiaire supérieur de Hongrie. – Acta Biol. Szeged. 32, 39–48.

Key words: Alginite, *Botryococcus braunii*, biopolymer structure.

-- i: In vitro destruction of the exine of recent palynomorphs I. – Acta Biol. Szeged. 32, 49–60.

Key words: Palynology, recent, biopolymer structure.

-- j: On the problems of the exine nomenclature. – Acta Biol. Szeged. 32, 205–206.

Key words: Palynology, exine nomenclature.

-- k: Explosion of pollen grains under the electron beam effect of the scanning electron microscope. – Acta Biol. Szeged. 32, 207–208.

Key words: Palynology, recent scanning effect.

-- l: A complex study of plant microfossils of oil shale by LM, TEM and thin layer chromatography. (In Hungarian, summary in English). – Bot. Közlem. 73, 25–32.

Key words: Alginite, biopolymer structure, thin layer chromatography.

HERNGREEN, G. F. W., FELDER, W. M., KEDVES, M.–MEESSEN, J. P. M. T.: Micropaleontology of the Maestrichtian in Borehole Bunde, The Netherlands. – Rev. Palaeobot. Palynol. 48, 1–70.

Key words: Palynology, fossil, n. fspp., Upper Cretaceous, The Netherlands.

HETÉNYI, M.–KEDVES, M.: Organic geochemical characterization of brown coals by thermal degradation and modified Rock-Eval method. – Acta Miner.-Petr. 28, 95–108.

Key words: Organic Geochemistry, Palynology, Eocene, Hungary.

KEDVES, M.–SZEDERKÉNYI, T.: Investigations on the microscopic plant remnants and the radioactive element contents of some mud samples of the Hungarian Plain. – Acta Biol. Szeged. 32, 209–211.

Key words: Organic microfossils, radioactive materials.

1987

-- a: Degradation of the sporoderm under natural and in vitro conditions. – XIV. International Botanical Congress, Berlin (West), Germany 24 July to 1 August 1987., Abstract, 595.

Key words: Palynology, recent, fossil, TEM, biopolymer structure.

-- b: Altérations des associations sporo-polliniques préquaternaires 25 ans après le premier traitement du sédiment. – Travaux et documents de géographie tropicale. X^e Symposium APLF "Palynologie, Ecologie, Paléoécologie", Talence, Résumés, 70.

-- c: LM and EM studies on pollen grains of recent *Welwitschia mirabilis* HOOK. and *Ephedra* species. – Acta Bot. Hung. 33, 81–103.

Key words: Palynology, recent, LM, TEM, SEM, *Welwitschia*, *Ephedra*.

-- d: Methodological problems in the investigation of the biopolymer units of the sporoderm. – 20th Annual Meeting AASP. Programme and Abstracts, Halifax, Nova Scotia, Canada, 100.

Key words: Palynology, fossil, biopolymer structure.

-- e: Paleophytogeography of the *angiosperm* pollen grains during the Upper Cretaceous and the Tertiary I. – Acta Biol. Szeged. 33, 35–47.

Key words: Palynology, Paleophytogeography, Cretaceous-Tertiary.

-- f: In vitro destruction of the exine of recent palynomorphs II. – Acta Biol. Szeged. 33, 49–56.

Key words: Palynology, recent, *Taxus baccata*, partial degradation.

-- g: Molecular structures from the organic remnants of the carbonate manganese ore layers of the III. Shaft of Úrkút, Hungary. – Acta Biol. Szeged. 33, 57–62.

Key words: Palynology, fossil, biopolymer structure, Jurassic, Úrkút, Hungary.

-- h: Higher organized sporopollenin biopolymer structures and the explosion of the pollen grains under scanning effect. – Acta Biol. Szeged. 33, 163–165.

Key words: Palynology, biopolymer structure, scanning effect.

- i: *Dégradation expérimentale de la paroi pollinique*. – Actas de Palinologia (Actas del VI. Simposio de Palinologia, A.P.L.E.) Salamanca, septiembre de 1986, 395–408.
Key words: Palynology, fossil, biopolymer structure, Paleocene, Menat, France.

1987–1988

- a: Beszámoló az A.P.L.F. IX., a trópusi környezet palinológiája témakörben megrendezett szimpóziumáról (Montpellier, 1985. október 1–3). – Bot. Közlem. 74–75, 247–249.
Key words: Palynology, A.P.L.F. Symposium, review.
 -- b: Beszámoló az A.P.L.E. VI. szimpóziumáról (Salamanca, 1986. szeptember 24–26.). – Bot. Közlem. 74–75, 251–253.
Key words: Palynology, A.P.L.E. Symposium, review.

Book reviews

- a: MROZINSKA, J.: *Chlorophyta VI. Oedogoniophyceae: Oedogoniales*. In: Süßwasserflora von Mitteleuropa 14. (eds. ETTL, H., GERLOFF, J., HEYNIG, H.–MOLLENHAUER, D.). – VEB Gustav Fisher Verlag, Jena, 1985. – Bot. Közlem. 74–75, 152.
 -- b: NAGY, E.: A magyarországi Neogén sporomorfái. – Geologica Hungarica 47, Budapest. – Bot. Közlem. 74–75, 110.

1988

- a: Evolution and Biology of the *Normapolles*. – Evolution, systematics and fossil history of the *Hamaelidae*. An International Symposium University of Reading 22nd–25th March 1988, Abstract, 16.
Key words: Palynology, fossil, *Angiospermatophyta*, *Normapolles*, evolution.
 -- b: Quasi-crystalloid basic molecular structure of the sporoderm. – 7 International Palynological Congress Brisbane, Abstracts, 82.
Key words: Palynology, recent, quasi-crystalloid, biopolymer structure.
 -- c: Trends and problems of the researches of fossil spores and pollen grains. – Annales Universitatis Scientiarum Budapestinensis de Rolando Eötvös Nominatae, Sect. Geol. 28, 169–175.
Key words: Palynology, fossil, ultrastructure, phylogeny.
 -- d: Degrees of biopolymer organization of the sporoderm as a contribution to the new concept of Global Geosphere – Biosphere modelling. – 21st Annual Meeting A.A.S.P. Program and Abstracts, Houston, Texas, USA.
Key words: Palynology, fossil, quasi-crystalloid, biopolymer structure.
 -- e: Methodological problems in the investigation of the biopolymer units of the sporoderm. – Palynology 12, 242.
Key words: Palynology, recent, fossil, biopolymer structure, methods.
 -- f: Alterations des associations sporo-polliniques éocènes à la suite du traitement palynologique. – Revista Española de Micropaleontología 20, 389–399.
Key words: Palynology, fossil, Eocene, Hungary, secondary selective fossilization.
 -- g: Paleophytogeography of the *angiosperm* pollen grains during the Upper Cretaceous and the Tertiary II. – Acta Biol. Szeged. 34, 45–57.
Key words: Palynology, Paleophytogeography, Cretaceous-Tertiary.
 -- h: Degradation of the sporoderm under natural and in vitro conditions. – Acta Biol. Szeged. 34, 59–69.
Key words: Palynology, recent, fossil, biopolymer structure.
 -- i: About the symmetry of the pentagonal basic biopolymer units of the pollen wall. – Acta Biol. Szeged. 34, 157–159.
Key words: Palynology, recent, quasi-crystalloid, biopolymer structure.
 -- j: First observation on the higher organized biopolymer structures of the exine of bisaccate *gymnosperm* pollen grains. – Acta Biol. Szeged. 34, 161–163.
Key words: Palynology, recent, biopolymer structure.

ABOUL ELA, N. M.–KEDVES, M.: Palynological studies on the intercalated sediments of the Yemen volcanics near Sana'a. – *Annales Universitatis Scientiarum Budapestinensis de Rolando Eötvös Nominatae, Sect. Geol.* 28, 27–41.

Key words: Palynology, fossil, reworking, Upper Cretaceous – Tertiary, Yemen.

KEDVES, M. J., KINCSEK, I., AMBRUS, E., FEJES, A.–GYEBROVSZKI, B.: La estructura molecular de la exina en algunos granos de polen bialados de *gymnospermas*. – *Polen, esporas y sus aplicaciones. VII Simposio de Palinologia. Granada, 26/30 Septbre. 1988.*, Resúmenes, 67.

Key words: Palynology, *Gymnospermatophyta*, biopolymer structure.

KEDVES, M.–SZEDERKÉNYI, T. (1988): Transmission electron microscopical investigation of xylem remains transporting radioactive elements in the mud of lake Vadkert. – *Acta Biol. Szeged.* 34, 71–81.

Key words: Palynology, Xylotomy, ultrastructure, radioactive elements transport.

KEDVES, M.–WINTER, J. (1988): Higher organized sporoderm biopolymer units of *Equisetum arvense* L. – *Acta Bot. Hung.* 34, 361–374.

Key words: Palynology, recent, *Equisetum arvense*, biopolymer structure.

NAGY, E.–KEDVES, M. (1988): State of palynological research in Hungary. – *Acta Bot. Hung.* 34, 311–324.

Key words: Palynology, research fields, review, Hungary.

1989

-- a: Palynologie et Paléocologie du Maestrichtien de l'Egypte. – First Symposium of African Palynology. Rabat, 15–21. May 1989. Abstracts.

Key words: Palynology, fossil, Upper Cretaceous, Egypt.

-- b: Transmission electron microscopical investigations on partially degraded plant cell walls. – Vth Symposium of the Hungarian Plant Anatomy. Szeged 25–26 August 1989, 22.

Key words: Plant cell wall, partial degradation, biopolymer structure.

-- c: Evolution of the *Normapolles* complex. In: Evolution, Systematics, and Fossil History of the *Hamamelidae* Volume 2 'Higher *Hamamelidae*', eds.: CRANE, P. R. and BLACKMORE, S., Clarendon Press Oxford, Systematics Association Special Volume 40B, 1–7.

Key words: Palynology, fossil, *Normapolles*, evolution.

-- d: Méthode d'étude des biopolymères de la paroi pollinique à structure quasi-cristalloïde. A method of investigation of the quasi-crystalloid structure of pollen wall biopolymers. – *Rev. de Micropaléontologie* 32, 226–234.

Key words: Palynology, recent, quasi-crystalloid biopolymer organization, methods.

-- e: New trends in micropaleontological researches. – II. European Palaeobotanical Conference, Madrid, September 1989, Abstracts, 3.

Key words: Palaeobotany, new methods, biopolymer organization.

-- f: Degrees of biopolymer organization of the sporoderm as a contribution to the new concept of Global Geosphere-Biosphere modelling. – *Palynology* 13, 284.

Key words: Palynology, biopolymer structure, Biosphere-Geosphere modelling.

-- g: Quasi-crystalloid biopolymer structures of the sporoderm and its highly organized degrees. – *Acta Biol. Szeged.* 35, 59–70.

Key words: Sporopollenin, quasi-crystalloid biopolymer structure, organizations model.

-- h: Beszámoló a XIV. Nemzetközi Botanikai Kongresszus palinológiai eredményeiről. – *Bot. Közlem.* 76, 147–149.

Key words: Botany, Palynology, XIV. I B. C., review.

-- i: Beszámoló az A.P.L.F. X. szimpóziumáról (Bordeaux, 1987, szept. 28 – okt. 2.). – *Bot. Közlem.* 76, 151–153.

Key words: Palynology, A.P.L.F. Symposium, review.

-- j: Beszámoló a "Hamamelidae" szimpóziumról (Evolution, Systematics and Fossil History of the Hamamelidae, Reading, 1988. március 22–25). – *Bot. Közlem.* 76, 155–157.

Key words: Palynology, *Hamamelidae* Symposium, review.

KEDVES, M.–KINCSEK, I. (1989a): Quasi-crystalloid biopolymer organization of the fossil spore and pollen wall. – II. European Palaeobotanical Conference, Madrid, September 1989, Abstracts, 16.

Key words: Palynology, fossil, quasi-crystalloid biopolymer structure.

KEDVES, M.–KINCSEK, I. (1989b): Effect of the high temperature on the morphological characteristic features of the sporomorphs I. – *Acta Biol. Szeged.* 35, 233–235.

Key words: Palynology, recent, *Amentiflorae*, high temperature effect.

KEDVES, M.–ROJIK, I. (1989): Investigations of the biopolymer organization of partially degraded exines with the fragmentation method. – *Acta Biol. Szeged.* 25, 71–80.

Key words: Palynology, sporopollenin, biopolymer organization, fragmentation method.

GÉVAY, G.–KEDVES, M. (1989): A structural model of the sporopollenin based on dodecahedrane units. – *Acta Biol. Szeged.* 35, 53–57.

Key words: Sporopollenin, biopolymer skeleton, structural model.

1990

-- a: Transmission electron microscopy of the fossil angiosperm exines. – Szeged, ISBN 963 481 8056.

Key words: Palynology, fossil, *Angiospermatophyta*, exine ultrastructure.

-- b: Palaeophytogeography of the Upper Cretaceous of the European and North African regions on the basis of palynological data. – International Symposium of the Nonmarine Cretaceous Correlation. International Geological Correlation Program Project No. 245, Alma-Ata 1990, Abstracts, 26–27.

Key words: Palynology, fossil, Upper Cretaceous, Paleophytogeography.

-- c: Experimental investigations on recent *Selaginella* spores. – *Taiwania* 35, 240–252.

Key words: Palynology, recent, *Selaginella*, LM, high temperature effect, TEM biopolymer structure.

-- d: Palynological studies on manganese ore layers in Úrkút (Transdanubia), Hungary. – *Ore Geology Reviews* 5, 491–507.

Key words: Palynology, fossil, Jurassic, LM, TEM, SEM, manganese ore, Úrkút, Hungary.

-- e: Quasi-crystalloid basic molecular structure of the sporoderm. – *Rev. Palaeobot. Palynol.* 64, 181–186.

Key words: Palynology, sporopollenin, quasi-crystalloid biopolymer structure.

-- f: Effect of the high temperature to the spores and pollen grains. – 8. Simposio de Palinologia A.P.L.E., Tenerife (Islas Canarias), 24/28. Septiembre 1990, Resúmenes, 43.

Key words: Palynology, recent, high temperature effect.

-- g: Highly organized biopolymer structures from the ectexine of *Thalictrum flavum* L. – *Acta Biol. Szeged.* 36, 99–101.

Key words: Palynology, recent, *Angiospermatophyta*, biopolymer structure.

KEDVES, M.–AILER, P. (1990): Investigations on recent monosulcate *Gymnospermatophyta* pollen grains. – *Acta Biol. Szeged.* 36, 103–105.

Key words: Palynology, recent, *Cycadales*, high temperature effect.

KEDVES, M., KINCSEK, I., AMBRUS, E., FEJES, A.–GYEBROVSZKI, B. (1990): Molecular structure of the exine of some *gymnospermous* bisaccate pollen grains. – In: *Polen esporas y sus aplicaciones*, eds: BLANCA, G., DIAZ DE LA GUARDIA, C., FERNÁNDEZ, M. C., GARRIDO, M., RODRIGUEZ-GARCIA, M. S. y ROMERO, A. T., Granada, 117–121.

Key words: Palynology, recent, *Gymnospermatophyta*, biopolymer structure.

HETÉNYI, M.–KEDVES, M. (1990): Relations between the hydrocarbon genetic features of kerogens and their biological precursor material. In: *Methods of Geochemical Prospecting Extended Abstracts*, eds: JANÁTKA, J., HLAVATÁ, T., BARNET, I. and JELÍNEK, E., Geological Survey, Prague, 246.

Key words: Organic Geochemistry, kerogens, biological precursor material.

1991

-- a: First observations on the biopolymer organization of the intine. – *Plant Cell Biology and Development* 1, 15–27.

Key words: Palynology, recent, *Gymnospermatophyta*, biopolymer structure.

-- b: Kvázi-kristalloid biopolymer struktúrák növényi sejtfalból. Quasi-crystalloid biopolymer structures from plant cell wall. – *Bio Társ* 7, 31.

Key words: Plant cell wall biopolymer organization.

-- c: Illustrations of the quasi-crystalloid biopolymer structures from the explosive dangerous coal pulver. – *Plant Cell Biology and Development* 2, 34–35.

Key words: Coal pulver, quasi-crystalloid biopolymer structures.

-- d: TICOS polyhedra as a model in the pentasporan organization. – Plant Cell Biology and Development 2, 43–48.

Key words: Palynology, recent, *Gymnospermatophyta*, biopolymer modelling.

-- e: Three dimensional modelling of the biopolymer structure of the plant cell wall I. – Plant Cell Biology and Development 2, 63–74.

Key words: Plant cell wall, biopolymer, three dimensional modelling.

-- f: Report on the activities of Palaeobotanical Subcommission (In Hungarian, summary in English). – Óslénytani Viták (Discussiones Palaeontologicae) 36–37, 67–76.

Key words: Hungarian Palaeobotany, review.

-- g: The biopolymer organization of the organic plant microfossils (In Hungarian, summary in English). – Óslénytani viták (Discussiones Palaeontologicae) 36–37, 77–89.

Key words: Plant cell wall, biopolymer organization.

-- h: Les modèles à trois dimension de l'organisation biopolymère du sporoderme. – Biogeographie et Palynologie, XII^e Symposium A.P.L.F. Caen 23–27, Septembre 1991, Résumés.

Key words: Sporoderm, biopolymer, three dimensional modelling.

KEDVES, M.–FARKAS, E. (1991): Basis of the tertiary rotation and TICOS modelling of the quasi-crystalloid biopolymer skeleton of the plant cell. – Plant Cell Biology and Development 2, 36–42.

Key words: Sporoderm, biopolymer, two dimensional modelling.

KEDVES, M., FARKAS, E., MÉSZÁROS, K., TÓTH, A.–VÉR, A. (1991): Investigations of the basic biopolymer structure of the ectexine of *Alnus glutinosa* (L.) GAERTN. – Plant Cell Biology and Development 2, 49–58.

Key words: Palynology, *Angiospermatophyta*, *Alnus*, biopolymer organization.

KEDVES, M., PÁRDUTZ, Á., FARKAS, E.–VÉR, A. (1991): Basic establishments of the biological objects molecular structure containing quasi-crystalloid skeleton. – Plant Cell Biology and Development 1, 35–37.

Key words: Quasi-crystalloid skeleton, basic establishments.

KEDVES, M., PÁRDUTZ, Á.–VÉR, A. (1991): Biopolymer organization of partially degraded exines of saccate *gymnosperm* pollen grains. – Plant Cell Biology and Development 1, 32–34.

Key words: Palynology, *Gymnospermatophyta*, biopolymer organization.

KEDVES, M.–ROJIK, I. (1991): Quasi-crystalloid biopolymer organization from the sclereids of *Armeniaca vulgaris* LAM. – Plant Cell Biology and Development 2, 59–62.

Key words: Sclereids, *Armeniaca vulgaris*, quasi-crystalloid biopolymer structure.

KEDVES, M., ROJIK, I.–VÉR, A. (1991): Biopolymer organization of the partially degraded oil shale with the fragmentation method. – Plant Cell Biology and Development 1, 28–31.

Key words: Oil shale, *Botryococcus braunii*, biopolymer organization.

KEDVES, M., TÓTH, A.–FARKAS, E. (1991): High temperature effect on the spores of *Equisetum arvense* L. – Plant Cell Biology and Development 1, 8–14.

Key words: Palynology, *Equisetum arvense*, high temperature effect.

KEDVES, M., TÓTH, A.–FARKAS, E. (1991): Effect of the high temperature on the morphological characteristic features of the sporomorphs II. – Acta Biol. Szeged. 37, 25–44.

Key words: Palynology, *Gymnospermatophyta*, monosulcate, inaperturate types, high temperature effect.

EL-SAADAWI, W. E.–KEDVES, M. (1991): Palaeobotanical investigations on plant impressions and sporomorphs from Egypt. – Plant Cell Biology and Development 2, 8–33.

Key words: Palaeobotany, macro- and microremnants, Lower Cretaceous, Egypt.

HETÉNYI, M.–KEDVES, M. (1991): Relations between the hydrocarbon genetic features of kerogens and their biological precursor material. – Exploration Geochemistry 1990, 128–132.

Key words: Organic Geochemistry, kerogens, biological precursor material.

1992

-- a: Three dimensional modelling of the biopolymer structure of the plant cell wall II. – Plant Cell Biology and Development 3, 67–87.

Key words: Plant cell wall, biopolymer, three dimensional modelling.

- b: Upper Cretaceous Paleophytogeography on palynological basis. – 8th International Palynological Congress, Aix-en-Provence, September 6–12, 1992, Abstracts, 74.
Key words: Palynology, Upper Cretaceous, Paleophytogeography.
- c: Études palynologiques des couches du Tertiaire inférieur de la Région Parisienne. VII. – Acta Biol. Szeged. 38, 33–46.
Key words: Palynology, *Angiospermatophyta*, Lower Tertiary, Paris Basin, France.
- d: Quasi-crystalloid biopolymer structures from the explosive dangerous coal pulver from Hungary. – Ann. Univ. Sci. Budapest., Sect. Geol. 29, 281–284.
Key words: Coal pulver, quasi-crystalloid biopolymer structures.
- e: Studies of the plant microfossil remnants of the Southern Bakony (In Hungarian, summary in English). – Óslénytani Viták (Discussiones Palaeontologicae) 38, 47–57.
Key words: Palynology, fossil, Eocene, Hungary.
- f: Biopolymer struktúrák szimmetriája. In: Szimmetria-Asszimmetria Megközelítések, Értelmezések – MTA SZTB, Szeged, ed.: BALOGH, T., 5–18.
Key words: Biopolymer structures, symmetry.
- KEDVES, M., FARKAS, E., GOTTL, E., MÉSZÁROS, K.–TÓTH A. (1992): L'importance des formes secondaires des sporomorphes produites par voie expérimentale. – 8th International Palynological Congress, Aix-en-Provence, September 6–12, 1992, Abstracts, 75.
Key words: Palynology, experimental secondary alterations.
- KEDVES, M., HEGEDŰS, A.–OLÁH, I. (1992): High temperature effect of some bisaccate *gymnosperm* pollen grains. – Plant Cell Biology and Development 3, 14–37.
Key words: Palynology, recent, *Gymnospermatophyta*, high temperature effect.
- KEDVES, M.–PÁRDUTZ, Á. (1992a): Transmission electron microscopy of partially dissolved exine of different bisaccate *gymnosperm* pollen grains. – Plant Cell Biology and Development 3, 38–66.
Key words: Palynology, *Gymnospermatophyta*, biopolymer structure.
- KEDVES, M.–PÁRDUTZ, Á. (1992b): TEM study of ultrathin sections of the partially degraded wall of the sclereids of *Armeniaca vulgaris* LAM. – Plant Cell Biology and Development 3, 88–91.
Key words: Sclereids, *Armeniaca vulgaris*, biopolymer structure.
- KEDVES, M.–PÁRDUTZ, Á. (1992c): X-ray effect to the exine ultrastructure of *Alnus glutinosa* (L.) GAERTN. – Taiwanica 37, 123–133.
Key words: Palynology, recent, *Alnus*, exine ultrastructure, X-ray effect.
- KEDVES, M., ROJIK, I.–VÉR, A. (1992): Ultrastructure and biopolymer organization of the *Botryococcus* colonies from Hungarian alginite. – Workshop on Pyrolysis in Organic Geochemistry. International Workshop, June 9–11, Szeged, 1992, Abstract, 21–22.
Key words: Oil shale, *Botryococcus braunii*, biopolymer structure.
- KEDVES, M.–TÓTH, A. (1992): Premiers résultats du système de biopolymère stabilisateur du squelette quasi-cristalloïde de l'exine. – 9. Simposio de Palinologia A.P.L.E., Las Palmas de Gran Canaria, Islas Canarias, 30 Noviembre al 4 Diciembre de 1992, Resúmenes, 20.
Key words: Palynology recent, biopolymer structure, stabilizing molecular system.
- KEDVES, M. TÓTH, A.–FARKAS, E. (1992): Experimental investigation of the biopolymer organization of the sporoderm (recent and fossil). – 8th International Palynological Congress, Aix-en-Provence, September 6–12, 1992, Abstracts, 75.
Key words: Sporoderm, biopolymer structure, symmetry.
- KEDVES, M., TÓTH, A., FARKAS, E., BELLON, A.–SCHMÉL, Á. (1992): Methodical problems of the biopolymer organization of partially degraded ectexine. – Ann. Univ. Sci. Budapest., Sect. Geol. 29, 263–279.
Key words: Sporoderm biopolymer two dimensional modelling.
- ALVAREZ RAMIS, C., KEDVES, M.–FERNÁNDEZ MARRÓN, T. (1992): Asociaciones esporopolínicas del Cretácico superior del Cerro de la Mesa (Guadalupe de la Sierra, Provincia de Madrid). – 9. Simposio de Palinologia A.P.L.E., Las Palmas de Gran Canaria, Islas Canarias, 30 Noviembre al 4 Diciembre de 1992, Resúmenes, 28.
Key words: Palynology, fossil, Upper Cretaceous, Spain.
- FERNÁNDEZ MARRÓN, M. T.–KEDVES, M. (1992): Sobre el carácter interdisciplinar de los estudios paleobotánicos. – Plant Cell Biology and Development 3, 11–13.
Key words: Paleobotany, interdisciplinary character.
- JELÉN, B., ANICIC, B., BREZIGAR, A., BUSER, S., CIMERMAN, F., DROBNE, K., MONOSTORI, M., KEDVES, M., PAVSIC, J.–SKABERNE, D. (1992): Model of positional relationships for Upper Paleogene and Miocene strata

in Slovenia. – I.U.G.S.–S.C.G. Miocene Columbus Projekt, Portonovo (Ancona, Italy), 11–14 November, 1992, Abstracts, 71–72.

Key words: Biostratigraphy, Tertiary, Slovenia.

SAJGÓ, Cs., HETÉNYI, M.–KEDVES, M. (1992): Palynology and organic geochemistry of Tertiary low rank coals in Hungary. – Workshop on Pyrolysis in Organic Geochemistry. International Workshop, June 9–11, Szeged, 1992, Abstracts, 36–37.

Key words: Organic Geochemistry, kerogens, biological precursor material.



A photograph of Prof. Dr. C. ALVAREZ RAMIS in the office of Dr. M. KEDVES.
The picture was taken by Dr. É. SIPOS-KEDVES.

Chronicle

Compiled by

M. KEDVES and I. GÁSPÁR

Visiting scientists

Dr. Bogomir JELEN, Senior Research Associate (Geological Survey of Ljubljana, Institute for Geology, Geotechnics and Geophysics, Ljubljana, Republic of Slovenia), visited the laboratory from 18th to 23th January, 1993. The aim of his visit was to work out the partial results of the joint research program dealing with the following Paleogene layers: Sava Folds (Zagorje-Lacko Syncline), Lower "Socka Beds", coal, Upper "Socka Beds", Oligocene Marine Clay.

Prof. Dr. C. ALVAREZ RAMIS (Department of Paleontology, Faculty of Geology, U.C.M. Madrid, Spain), spent two weeks in Szeged from 28th May to 11th June, 1993 and worked on the preparation of the publication of the results of the joint research program: "Estudio de diversos aspectos paleobotánicos del Cretácico superior del Cerro de la Mesa, Norte de la provincia Madrid."

On the 1st June, 1993 she took part in the scientific session organized in Budapest by the Biological Department of the Hungarian Academy of Sciences on the occasion of the 85th birthday of Prof. Dr. B. ZÓLYOMI academician, establisher of the Hungarian Quaternary Palynology. Prof. Dr. M. KEDVES and A. VÉR represented the laboratory on this event.

Dr. Bogomir JELEN visited again the laboratory in December (13–17, 12, 1993) and worked out the first part of the previously mentioned program for publication.

International laboratory activity

20–22 May, 1993 – Berlin

Scientific meeting dedicated at the occasion of Prof. Dr. W. KRUTZSCH's 65th birthday, organized by Prof. Dr. R. DABER (Museum für Naturkunde, Berlin) and Dr. H. BLUMENSTENGEL (Geologisches Landesamt Sachsen-Anhalt, Berlin).

KEDVES, M.: Multidisciplinary characters of Palaeopalynological researches.

27 August–5 September, 1993 – Yokohama

XVth International Botanical Congress, Yokohama, Japan. Dr. M. KEDVES took part in the congress as an invited speaker in the section S1.8.1., "The Contribution of Pollen and Spore Studies to Systematic and Evolutionary Biology (Stephen BLACKMORE, Masamichi TAKAHASHI);

KEDVES, M., TÓTH, A. and VÉR, A.: Radial Fivefold Rotation: A New Method in the Study of the Biopolymer Organization of the Sporoderm.

18–26 September, 1993 – Besançon

The XIIIth Symposium of the A.P.L.F. "Palynologie et événements cycliques" Besançon, 20–24 September, 1993, France.

KEDVES, M. et BAGI, I.: Études palynologiques et pédologiques sur les sédiments Holocènes de la piscine du monastère du Mont Jakab.

11–25 October, 1993 – Madrid.

Prof. Dr. M. KEDVES worked in the Department of Paleontology (Facultad de Ciencias, Universidad Complutense de Madrid) in Madrid as a visiting scientist together with Prof. Dr. C. ALVAREZ RAMIS and Dr. M. T. FERNÁNDEZ MARRÓN. They closed the previously mentioned joint research program, considered the final results and prepared a new joint research program.

Laboratory meetings

5. 2. 1993, speaker: KEDVES, M.: Discussion of the up-to-date research programs of the laboratory.
2. 4. 1993, 1. speaker: GÁSPÁR, I.: Review of two papers written about recent and fossil *Equisetum* spores.
 2. speaker: VÉR, A.: Discussion of her certificate dissertation.
 3. speaker: KEDVES, M.: Actual problems.
- 12.11.1993, speaker: KEDVES, M.: Report on the XIIIth Symposium of the A.P.L.F. "Palynologie et événements cycliques". 20–24 September, 1993, Besançon, France.
22. 12. 1993, speaker: KEDVES, M.: Discussion of the participation of the laboratory on the international scientific meetings in 1994.

Preparation of the next number of the laboratory bulletin.

Teaching program of the laboratory

During the two semesters in 1993 the following lectures were delivered:
Introduction to the plant micropaleontology of pre-Quaternary deposits, 1+2 hours weekly.

Organization levels of the biopolymer system of the plant cell wall, 1+2 hours weekly.
Theory of the evolution and its natural philosophical relations, 1 lecture weekly.

Personalia

Anita TÓTH is one of the establishers of our laboratory. She has worked here with a great diligence and persistently since she was a grammar-school student. Later she attended to the Szentgyörgyi Albert Medical University in Szeged and now she is in her fourth year. On 3rd October, 1992 she married and gave birth to her daughter, Viola on 2nd April, 1993.

B 14 6191



Responsible for publication: M. KEDVES
Responsible editor: Zs. PAPP
Cover by E. KOVÁCS
Set in New Times 10/11 point
Juhász Nyomda, Szeged

